

NO LOCAL DOUBLE EXPONENTIAL GRADIENT GROWTH IN HYPERBOLIC FLOW FOR THE 2D EULER EQUATION

VU HOANG AND MARIA RADOSZ

ABSTRACT. We consider smooth, double-odd solutions of the two-dimensional Euler equation in $[-1, 1]^2$ with periodic boundary conditions. This situation is a possible candidate to exhibit strong gradient growth near the origin. We analyze the flow in a small box around the origin in a strongly hyperbolic regime and prove that the compression of the fluid induced by the hyperbolic flow alone is not sufficient to create double-exponential growth of the gradient.

1. INTRODUCTION

The question whether solutions of the two-dimensional Euler equation in vorticity form

$$(1) \quad \omega_t + u \cdot \nabla \omega = 0, \quad u = \nabla^\perp (-\Delta)^{-1} \omega$$

($\nabla^\perp = (-\partial_{x_2}, \partial_{x_1})$) can exhibit strong gradient growth in time is a topic of ongoing interest. The best known upper bound predicts double-exponential growth in time:

$$(2) \quad \|\nabla \omega(\cdot, t)\|_{L^\infty(\Omega)} \leq C_1 \exp(C_2 \exp(C_3 t))$$

on a domain Ω with either a smooth boundary with no-flow boundary condition or no boundary (e.g. a torus). The constants C_i depend on the initial data. A natural and important question is: Are there flows for which this upper bound is attained? For domains with boundary, a recent breakthrough by A. Kiselev and V. Šverák [8] answers the question affirmatively. In [8], solutions are constructed that attain the double-exponential bound (2).

For smooth solutions on the torus, the situation is far from clear. The best known result so far was given by S. Denisov. In [4], he shows that at least superlinear gradient growth is possible and in [5] he provides an example of double-exponential growth for an arbitrarily long, but finite time interval. In the recent paper [11], A. Zlatoš constructs initial data leading to exponential gradient growth, his solution is however in $C^{1,\gamma}$ for some $\gamma \in (0, 1)$ and not in C^2 .

In [8] the construction is based on creating a *hyperbolic flow scenario*. By imposing a symmetry on the solutions, a stagnant point of the flow is created on the boundary of the domain. The initial conditions are chosen in such a way the flow on the boundary is directed towards the stagnant point, creating a strong fluid compression and therefore strong gradient growth.

A natural way to carry the Kiselev-Šverák construction to the torus is to consider double-odd solutions, i.e.

$$(3) \quad \omega(-x_1, x_2) = -\omega(x_1, x_2), \quad \omega(x_1, -x_2) = -\omega(x_1, x_2),$$

Date: January 19, 2016.

This construction was employed in [11]. In [5], a perturbation argument starting from a non-smooth double-odd stationary solution (see [1]) was used. So far, however, creating infinite-time *double-exponential* growth in the double-odd scenario was not successful. Our goal in this paper is to explore the difficulties in using this scenario, by proving a conditional regularity result.

It is interesting to notice that the result [8] is in some sense analogous to the still open blowup problem for the more singular surface quasigeostrophic equation. In SQG blowup means that the solution becomes singular in finite time whereas for the 2d Euler equation “blowup” would mean maximal (double-exponential) gradient growth on an infinite time interval. There are important conditional regularity results for the SQG equation such as [2, 3], where the authors study a certain blowup scenario, in order to finally exclude it. An analogous “conditional regularity result” for 2d Euler equation would be to show that in certain scenarios maximal gradient growth does not occur. Since the possible motions of fluids are various and in general very complicated, studying scenarios is an invaluable method to gain insight into regularity problems of fluid mechanics.

Our main result states that a hyperbolic flow cannot create double-exponential gradient growth near the origin by itself when we start with double-odd C^2 initial data, provided a certain “upstream” control is assumed on the flow. This is an important step into understanding the double-odd hyperbolic scenario since we rule out the most promising candidate for a mechanism creating maximal gradient growth, i.e. the local hyperbolic compression. Our result does not imply impossibility of double-exponential growth in general, but makes the construction of examples much harder.

In some sense, the scenario considered here is complementary to the one considered by D. Cordoba for the SQG equation in [2], where a closing hyperbolic saddle is considered. There the solution stays smooth except for the possible closing of the saddle. In our scenario for 2d Euler, the hyperbolic saddle is fixed due to the symmetry ($\omega = 0$ on the coordinate axes), and we are asking if blowup can happen in another way.

We strongly believe that the techniques developed here will also be useful in understanding the hyperbolic scenario for other models in fluid mechanics and also in situations with a physical boundary. There, although the goal is to prove the existence of a blowup, a certain amount of control up to the blowup time is necessary.

Interesting results concerning the related question of existence of double-exponential growth in the context of (nonsmooth) patch solutions were given by S. Denisov (see [6]).

Finally, we would like to mention the recent preprint [7], where a different approach is proposed to study whether double-exponential gradient growth can occur at an interior point.

1.1. Setup and feeding conditions. We consider (1) on $\mathbb{T} = [-1, 1]^2$ with periodic boundary conditions and double-odd C^2 initial data ω_0 . From now on, we use $\|\cdot\|_\infty$ to denote the L^∞ -norm on the torus $\mathbb{T}$.

The double-odd symmetry is preserved by the evolution and (3) implies that the origin is a stagnant point of the flow field for all times. Moreover, the flow on each coordinate axis is always directed along that axis. When considering smooth solutions $\omega \in C^1([0, \infty), C^2(\mathbb{T}))$, (3) also implies

$$\omega = 0$$

on the coordinate axes.

We will study the flow in boxes of the form

$$D = (0, \delta_1) \times (0, \delta_2), \quad \widehat{D} = (0, \delta_1 + \delta_3) \times (0, \delta_2),$$

where δ_j are positive, but small and

$$0 < \delta_1 < \delta_2 < \delta_1 + \delta_3.$$

In a hyperbolic flow, the origin is a stagnant point of the flow and fluid particles constantly enter the box D from the right and leave on the top (see Fig. 1). The particles moving on the x_1 -axis approach asymptotically the origin and never leave the box D . Generally speaking, there is, a compression of the fluid in x_1 -direction and a decompression in x_2 -direction (or the other way around). The vorticity is zero on the axes. The gradient growth in the box D comes from two sources: particles that were at $t = 0$ inside D and those which enter the box at later times. The time evolution of the gradient of those particles entering the box is difficult to control over infinite times, and is generated by flow situations which have little to do with the hyperbolic scenario. We are interested in making local statements and must *assume* a certain control on the flow entering the box D .

We shall therefore call $\widehat{D} \setminus D$ *feeding zone* and formalize this idea in the following definition (the meaning of the parameter α will become clear later).

Definition 1.1. Let $\alpha \in (0, \frac{1}{4})$. The box $\widehat{D}$ is said to satisfy the conditions of *controlled feeding*, with feeding parameter $R \geq 0$ if

$$(4) \quad \left| \frac{\partial \omega}{\partial x_1}(x, t) \right| \leq R x_2^{1-\alpha}, \quad \left| \frac{\partial \omega}{\partial x_2}(x, t) \right| \leq R \quad (x \in \widehat{D} \setminus D)$$

for all times $t \geq 0$.

We can think of the first inequality in (4) as a Hölder-version of a bound on $\partial_{x_2, x_1} \omega$, keeping in mind that $\partial_{x_1} \omega(x_1, 0, t) = 0$ for all times. The concept of controlled feeding conditions allows us to study the evolution of ω in D independent of the remaining flow. Note that for the purposes of this paper, we consider time-independent R only (see also Remark 2.1).

1.2. The hyperbolic scenario. In order to give a definition of hyperbolic flow suitable for our purposes, we introduce the following important quantity. Let $\alpha \in (0, \frac{1}{4})$ be fixed. For a smooth, periodic function ω we set

$$M(x, t) := \max_{0 \leq y_1, y_2 \leq \max\{x_1, x_2\}} \left\{ \left| y_1^\alpha \frac{\partial \omega}{\partial x_1}(y, t) \right|, \left| y_2^\alpha \frac{\partial \omega}{\partial x_2}(y, t) \right| \right\} + \|\omega\|_\infty.$$

Note that $M(x, t)$ also depends on ω and α . The velocity field $u(x, t) := \nabla^\perp(-\Delta)^{-1} \omega$ for double-odd ω (ω with mean zero over $\mathbb{T}$) can be written in the form

$$(5) \quad u_1(x, t) = -x_1 Q_1(x, t), \quad u_2(x, t) = x_2 Q_2(x, t)$$

where Q_1, Q_2 are scalar fields given by certain integral operators (see (14)) acting on ω . The following definition states that we regard the flow as hyperbolic if both Q_1 and Q_2 essentially have a positive lower bound, up to a term controlled by the quantity $M(x, t)$.

Definition 1.2. Let ω be a smooth solution of the Euler equation, and let $\alpha \in (0, \frac{1}{4})$ be fixed. We say that the flow is hyperbolic near the origin if there are constants $\rho, A, \beta_0 > 0$ for which the following condition is satisfied for all $t \in [0, \infty)$:

$$(6) \quad Q_i(x, t) + A|x|^{1-\alpha}M(x, t) \geq \beta_0 > 0 \quad (0 \leq x_1, x_2 \leq \rho, i = 1, 2).$$

The model situation for a hyperbolic flow is the following: Consider the dynamical system

$$\dot{x}_1 = -Ax_1, \quad \dot{x}_2 = Bx_2$$

with positive constants A, B . This system has a hyperbolic saddle point in $(0, 0)$, which is a stagnant point of the flow. On the axes, the flow is directed along the axes.

The velocity field given by (5) generalizes this structure if $Q_i(x, t) \geq \beta_0 > 0$. A further generalization necessary for our result is (6), since the stronger condition $Q_i(x, t) \geq \beta_0 > 0$ cannot be easily realized. In certain flow situations, the term $|x|^{1-\alpha}M(x, t)$ is small close to the origin. Bounding the quantity $M(x, t)$ plays a central role in our estimates.

Remark 1.3. By choosing the initial data ω_0 suitably, we can ensure hyperbolic flow. One possible choice is, for example, choosing ω_0 to be nonnegative in $[0, 1]^2$ and such that $\omega_0 = 1$ on a set of sufficiently large measure, as it was done in [8, 11]. This creates a situation where (6) is satisfied (see theorem 4.2). In this sense, (6) is a “realistic” condition on the flow.

1.3. Main result. Our main result is the following theorem.

Theorem 1.4. Fix $0 < \alpha < \frac{1}{4}, 0 < \delta_3 < 1/2$. Let ω be a C^2 , double-odd solution of the Euler equation with initial data ω_0 , and suppose the flow is hyperbolic near the origin. Let $R > \|\omega_0\|_\infty > 0$ be given. Then there exist small $\delta_1, \delta_2 > 0$ depending on α, β_0, R and ω_0 , such that if $\hat{D}$ satisfies the controlled feeding conditions with parameter R , then

$$\|\nabla \omega\|_{D, \infty} \leq C_1 \exp(C_2 t) \quad (t \in [0, \infty))$$

for some $C_1, C_2 > 0$ depending on $R, \alpha, \beta_0, \delta_1, \delta_2, \delta_3, \omega_0$.

This means that the hyperbolic compression alone and the interaction of the fluid inside the box is not sufficient to create double-exponential gradient growth. One would have to create a scenario where the feeding conditions are violated. This means roughly that there has to be a kind of compression in x_2 -direction in the feeding zone. This would have to be caused by much more complicated interactions outside the box. At the present time, no such scenario is known.

2. GRADIENT GROWTH IN THE HYPERBOLIC SCENARIO

Before describing our approach, let us explain first why at first sight the hyperbolic scenario seems to be a good candidate for double-exponential growth. Namely, for Q_1, Q_2 we have the upper bounds

$$Q_1(x, t), Q_2(x, t) \lesssim \|\omega\|_\infty |\log(x_1^2 + x_2^2)|.$$

If it were possible to create a situation where a *lower bound* of roughly the same order holds, i.e. $Q_1 \geq C |\log(x_1^2 + x_2^2)|$ over an infinitely long time interval, then for the particle trajectories lying on the x_1 -axis (i.e. $X_2 = 0$)

$$X_1(t) \leq \exp(-C_1 \exp(C_2 t))$$

would hold, as seen by solving the ODE $\dot{X}_1 = -X_1 Q_1$. If, moreover, one could arrange for the initial data ω_0 to have suitable nontrivial values on the x_1 -axis, then this would create double exponential gradient growth. However, the simultaneous requirements of smoothness and double-odd symmetry of ω , necessarily imply $\omega = 0$ on the axes. Moreover, it is highly unclear how a such strong lower bound on Q_1 could be achieved. As we shall see later, a certain amount of smoothness of ω and the vanishing of ω on the axes lead to a better upper bound, *without* the logarithmic behavior which is crucial for the double-exponential growth.

Another way one might hope to get double exponential growth is to consider a “projectile”, i.e. to track the movement of a small domain close to the origin on which $\omega = 1$, as it was done in [8]. There the self-interaction of the projectile was able to create enough growth in the values of Q_1 to allow double-exponential growth. While the projectile approaches the origin, the values of Q_1 on it get larger, this fact being connected to a certain logarithmically divergent integral. Our Theorem 1.4 shows that in general this is not possible for double-odd solutions, unless there is some kind of compression in x_2 -direction in the feeding zone. Thus a scenario with maximal gradient growth must be much more complicated than using the self-interaction of the projectile.

In fact, provided the feeding condition holds, the steady fluid compression guaranteed by (6) will turn out to stabilize the flow in the neighborhood of the origin. That is, the hyperbolicity condition (6) - which is essentially a lower bound on Q_i - is converted in the proof of Theorem 1.4 into an upper bound for Q_i . This is what finally leads to a bound on the gradient growth in D .

2.1. Heuristic considerations. We now present an intuitive discussion of our result. Fluid particles carried by the hyperbolic flow will constantly enter the box D from the right and leave on the top (see figure 1). All particles except for those moving on the axes spend a finite time in the box. The particles on the x_1 -axis move towards the left approaching the origin asymptotically as $t \rightarrow \infty$. Particle trajectories $t \mapsto \mathbf{X}(t) = (X_1(t), X_2(t))$ for which $X_2(0)$ is small approximate the straight trajectories of the particles on the x_1 -axis for a long time, before going steeply upward. The time a particle spends in D goes to infinity as $X_2(0) \rightarrow 0$.

We now consider the trajectory of a particle $\mathbf{X}$. The particle may have started inside D at time $t = 0$, or may have entered the box at some time $T_0 > 0$, in which case $\mathbf{X}(T_0) \in \partial D$. Also, assume that the particle exits the box D at some time T_e , i.e. $X_2(T_e) = \delta_2$. The evolution of the gradient of ω along the trajectory is given by an ODE of the form

$$(7) \quad \frac{d}{dt} \nabla \omega(\mathbf{X}(t), t) = (-\nabla u(\mathbf{X}(t), t))^T \nabla \omega(\mathbf{X}(t), t)$$

where ∇u is the velocity gradient. The relation (7) is simply derived by differentiating the Euler equation. The key is now to use the structure (5) of the velocity field such that we obtain

$$(8) \quad \frac{d}{dt} \nabla \omega(\mathbf{X}(t), t) = \begin{pmatrix} Q_1 + x_1 \frac{\partial Q_1}{\partial x_1} & -x_2 \frac{\partial Q_2}{\partial x_1} \\ x_1 \frac{\partial Q_1}{\partial x_2} & -Q_2 - x_2 \frac{\partial Q_2}{\partial x_2} \end{pmatrix} \nabla \omega(\mathbf{X}(t), t)$$




FIGURE 1. Illustration of a hyperbolic flow.

We write the right hand side of (8) as

$$\begin{pmatrix} a(t) & c(t) \\ b(t) & -a(t) \end{pmatrix} \nabla \omega(\mathbf{X}(t), t).$$

evaluating all matrix entries along the given trajectory $\mathbf{X}$. Note that the matrix has trace zero, since the velocity field u is divergence free. There are several ways we can heuristically regard (8) as a perturbation of an easier problem.

- For the discussion assume that $Q_1, Q_2 > 0$ and we can control the derivatives $\partial_{x_j} Q_i$ for small x . Since in a sufficiently small box $x_1 \partial_{x_1} Q_1, x_2 \partial_{x_2} Q_2$ should be rather “small” (due to the prefactors x_1, x_2), a should be positive and bounded away from zero along the hyperbolic trajectory. To gain some insight, we consider the case of a particle moving close to the x_1 -axis, i.e. with small $X_2(T_0) > 0$. We expect that $c = -x_2 \partial_{x_1} Q_2, b = x_1 \partial_{x_2} Q_1$ are “small”. Life would be easy if we could neglect b, c and set $b, c = 0$ in (8), so that we have a diagonal system. Denoting $\xi(t) = \nabla \omega(\mathbf{X}(t), t)$ the solution can be explicitly computed to be

$$(9) \quad \xi_1(t) = e^{A(t)} \xi_1(T_0), \quad \xi_2(t) = e^{-A(t)} \xi_2(T_0).$$

where $A(t) = \int_{T_0}^t a(\mathbf{X}(s)) ds$. (9) shows that, in general, the gradient in x_1 -direction grows along the particle trajectory. However, there is an effect which allows us to cancel the growing factor e^A . Assume for the sake of the discussion that the following stronger feeding conditions hold:

$$(10) \quad \left| \frac{\partial \omega}{\partial x_1}(x, t) \right| \leq R x_2, \quad \left| \frac{\partial \omega}{\partial x_2}(x, t) \right| \leq R,$$

on $\hat{D}$ for $t = 0$ and on $\hat{D} \setminus D$ for all $t > 0$. These imply in either case $T_0 = 0$ and $T_0 > 0$

$$(11) \quad |\xi_1(t)| \leq e^{A(t)} |\xi_1(T_0)| = e^{A(t)} \left| \frac{\partial \omega}{\partial x_1}(\mathbf{X}(T_0), T_0) \right| \leq R e^{A(t)} X_2(T_0).$$

Now we observe that

$$(12) \quad A(t) \approx \int_{T_0}^t Q_2(s) \, ds$$

temporarily neglecting the “small” term $x_2 \partial_{x_2} Q_2$. Now from (5) we have the differential equation $\dot{X}_2 = X_2 Q_2$, so that $X_2(t) = X_2(T_0) \exp \left(\int_{T_0}^t Q_2(\mathbf{X}(s)) \, ds \right)$ and hence

$$(13) \quad X_2(T_0) = X_2(T_e) \exp \left(- \int_{T_0}^{T_e} Q_2(\mathbf{X}(s)) \, ds \right) \leq \delta_2 \exp \left(- \int_{T_0}^{T_e} Q_2(\mathbf{X}(s)) \, ds \right).$$

Combining (13), (11) and (12), we get

$$|\xi_1(t)| \leq \delta_2 R \exp \left(- \int_t^{T_e} Q_2(\mathbf{X}(s)) \, ds \right) \leq \delta_2 R,$$

suggesting that the gradient in x_1 -direction *does not grow at all in time* given the feeding condition (10). Our rigorous result does not give such a strong conclusion, but we will be able to prove that the gradient *grows at most exponentially in time* using a weaker feeding condition. In Remark 4.6 we explain why (10) is not an appropriate feeding condition for the problem.

The heuristics appear deceptively simple, but in order to make the argument rigorous, we have to overcome a number of formidable technical difficulties. To begin with, the coefficients of (8) depend on the solution ω through the integral operators Q_1, Q_2 . The derivatives $\partial_{x_1} Q_1, \partial_{x_2} Q_2$ are given by singular integral operators.

Of course, none of the coefficients may be neglected, and we have to produce sufficiently good estimates on the solutions of the full ODE system (8). A major obstacle in getting good estimates, however, is caused by the unstable nature of (8). To illustrate this we consider a tridiagonal system by setting $c = 0$, but keeping b , so that we get a supposedly better approximation than the diagonal system. In this model, too, the solutions can be calculated explicitly, and we get

$$\xi_1(t) = e^{A(t)} \xi_1(T_0), \quad \xi_2(t) = e^{-A(t)} \left[\xi_2(T_0) + \xi_1(T_0) \int_0^t b(s) e^{2A(s)} \, ds \right].$$

This shows that not only the derivative in x_1 -direction but also the derivative in x_2 -direction of ω may potentially grow in time (due to the contribution $e^{-A(t)} \int_0^t b(s) e^{2A(s)} \, ds$). To make things worse, a possible strong growth in $\partial_{x_2} \omega$ is coupled back into the coefficients of the ODE (8) via our estimates on $\partial_{x_1} Q_1, \partial_{x_2} Q_2$. On the other hand, by a similar argumentation as in the case of the diagonal system, the factor $\xi_1(T_0)$ may help via a feeding condition. We need therefore to proceed with extreme care, looking to cancel the growing factor e^A with the decaying factor e^{-A} whenever possible.

Remark 2.1. In our scenario, we always assume the intensity of the feeding (i.e. the quantity R) to be time-independent. One might think of allowing the feeding parameter to grow in time to include more complicated scenarios. However, this is met with considerable challenges.

Firstly, it is not clear what a realistic condition on R should be, since it depends on the complexity of the flow away from the origin. One concrete situation where we can imagine a reasonable time-dependent feeding condition is as follows: A vortex created by a large

patch (see Figure 2) where ω is constantly 1. The flow revolves in clock-wise direction around the patch and, in analogy with a shear flow, one could assume linear growth in time of the gradient in the feeding zone.

The application of the techniques developed here to time-dependent feeding are not straightforward (see Remark 7.3), due to the non-local and non-linear nature of the problem.




FIGURE 2. Flow around a patch.

3. NOTATION

3.1. Euler velocity field. For $x = (x_1, x_2)$ we write $\tilde{x} = (-x_1, x_2)$ and $\bar{x} = (x_1, -x_2)$. The velocity field for the Euler equation is

$$u(x, t) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{(y - x)^\perp}{|y - x|^2} \omega(y, t) dy$$

where $\omega \in C^2(\mathbb{T})$ is periodically extended to all of $\mathbb{R}^2$ and $z^\perp = (-z_2, z_1)$. In the calculation of the integral a limit in the mean (sequence of unboundedly growing domains) is understood. Note that the velocity field is $\nabla^\perp(-\Delta)^{-1}\omega$, where $-\Delta$ is the periodic Laplacian on the Torus $\mathbb{T}$. A simple calculation using the double-odd symmetry of ω leads to

$$u_1(x, t) = -x_1 Q_1(x, t), \quad u_2(x, t) = x_2 Q_2(x, t)$$

where Q_1, Q_2 are the following integral operators (see Appendix B)

$$(14) \quad \begin{aligned} Q_1(x, t) &= c_0 \int_{[0,1]^2} [G_1^1(x, y) + G_1^2(x, y)] \omega(y, t) dy + Q_1^r(x, t) \\ Q_2(x, t) &= c_0 \int_{[0,1]^2} [G_2^1(x, y) + G_2^2(x, y)] \omega(y, t) dy + Q_2^r(x, t) \end{aligned}$$

with kernels

$$\begin{aligned} G_1^1(x, y) &= \frac{y_1(y_2 - x_2)}{|y - x|^2 |y - \tilde{x}|^2}, & G_1^2(x, y) &= \frac{y_1(y_2 + x_2)}{|y + x|^2 |y - \tilde{x}|^2}, \\ G_2^1(x, y) &= \frac{y_2(y_1 + x_1)}{|y + x|^2 |y - \tilde{x}|^2}, & G_2^2(x, y) &= \frac{y_2(y_1 - x_1)}{|y - x|^2 |y - \tilde{x}|^2}, \end{aligned}$$

where c_0 denotes the right constant. The expression Q_1^r is given by the following (limit in the mean) integral

$$c_0 \int_{\mathbb{R}_+^2 \setminus [0,1]^2} [G_1^1(x, y) + G_1^2(x, y)] \omega(y) dy, \quad \mathbb{R}_+^2 = (0, \infty)^2,$$

a similar formula holding for Q_2^r .

In section 4 we will derive estimates for the entries of the matrix in (8) which are independent of the trajectory. For this purpose it is convenient to use the definitions:

$$(15) \quad \begin{aligned} a(x, t) &:= Q_1(x, t) + x_1 \frac{\partial Q_1}{\partial x_1}(x, t) = Q_2(x, t) + x_2 \frac{\partial Q_2}{\partial x_2}(x, t), \\ b(x, t) &:= x_1 \frac{\partial Q_1}{\partial x_2}(x, t), \\ c(x, t) &:= -x_2 \frac{\partial Q_2}{\partial x_1}(x, t). \end{aligned}$$

Moreover, since the estimates will be for fixed t we shall often skip the t variable in the notation. When evaluating a, b, c, Q_i etc. along a particle trajectory $\mathbf{X}(t)$ in section 6 we shall write $a(t) := a(\mathbf{X}(t), t)$ etc. reconciling with the notation in section 2.1.

3.2. Convention for estimates. The notation $f \lesssim g$ means

$$f \leq Cg,$$

where C may depend on $\alpha, \beta, \|\omega\|_\infty$ and on universal constants, e.g. geometrical characteristics of the domain $\mathbb{T}$. C does not depend on $\delta_1, \delta_2, \delta_3, t$. When using this notation, we shall always imply that $C < \infty$ for all $\alpha \in (0, \frac{1}{4})$.

4. POTENTIAL THEORY OF Q_1, Q_2

4.1. Sufficient conditions for hyperbolic flow. We will be working with boxes of the form

$$(16) \quad \begin{aligned} D &= (0, \delta_1) \times (0, \delta_2) \\ \widehat{D} &= (0, \delta_1 + \delta_3) \times (0, \delta_2), \end{aligned}$$

with the following restriction:

$$(17) \quad 0 < \delta_1 < \delta_2 < \delta_1 + \delta_3.$$

and δ_j so small that $\widehat{D} \subset [0, 1]^2$. We also write

$$d(x) = \delta_2 - x_2$$

which is the distance of the point x to the top of the box. We write $\boldsymbol{\delta} = (\delta_1, \delta_2)$, $|\boldsymbol{\delta}|^2 = \delta_1^2 + \delta_2^2$.

Define

$$M_D(t) := \max_{y \in D} \left\{ \left| y_1^\alpha \frac{\partial \omega}{\partial x_1}(y, t) \right|, \left| y_2^\alpha \frac{\partial \omega}{\partial x_2}(y, t) \right| \right\} + \|\omega\|_\infty$$

and $M_{\widehat{D}}$ for the analogous quantity. Note that M_D and $M_{\widehat{D}}$ depend on ω and α .

As mentioned before, the flow near the origin can be made hyperbolic, with compression in the x_1 -direction and expansion in x_2 -direction by choosing the initial data such that $\omega_0 \geq 0$ on $[0, 1]^2$ and such that

$$\mathfrak{m} := |\{x : \omega_0(x) = \|\omega_0\|_\infty\}|$$

is sufficiently large. This is a consequence of theorem 4.2.

Remark 4.1. (a) As a consequence of $\omega = 0$ on the coordinate axes we have the following important inequality

$$(18) \quad |\omega(y, t)| \lesssim M(x, t) y_j^{1-\alpha} \quad (y_1, y_2 \leq \max\{x_1, x_2\})$$

where $j = 1, 2$.

(b) The periodicity and double-oddness of $\omega(\cdot, t)$ imply also the reflection symmetries

$$\omega(1 + x_1, x_2, t) = -\omega(1 - x_1, x_2, t), \quad \omega(x_1, 1 + x_2) = -\omega(x_1, 1 - x_2).$$

Consequently, the four corner points of $[-1, 1] \times [-1, 1]$ are also stagnant points of the flow, the flow being confined in $[0, 1]^2$. Hence $\omega_0 \geq 0$ on $[0, 1]^2$ implies $\omega(x, t) \geq 0$ on $[0, 1]^2$ for all times, a fact we shall use below.

Theorem 4.2. *Suppose $\omega_0(x) \geq 0$ on $[0, 1]^2$. There exist universal $0 < m_0 < 1$ and $0 < K$ such that if $m_0 < \mathfrak{m} < 1$, there are $\beta_0 > 0, A > 0$ such that the following estimates hold for all times*

$$(19) \quad \begin{aligned} Q_2(x, t) + AM(x, t)|x|^{1-\alpha} &\geq \beta_0 \\ Q_1(x, t) + AM(x, t)|x|^{1-\alpha} &\geq \beta_0 \end{aligned}$$

for $|x| \leq K(1 - \mathfrak{m})$, i.e. the flow is hyperbolic near the origin.

To prove this, we need the following Lemma, which is an adaption of a result in [11].

Lemma 4.3. *Let $\Omega(2x) := [2x_1, 1] \times [2x_2, 1]$. Suppose $\omega(x) \geq 0$ for $x \in [0, 1]^2$. Then the estimate*

$$Q_i(x, t) \geq c_0 \int_{\Omega(2x)} \frac{y_1 y_2}{|y|^4} \omega(y, t) dy - C_1 M(x, t) |x|^{1-\alpha} - C_2 \|\omega\|_\infty \quad (x \in D, i = 1, 2)$$

holds, with universal $C_1, C_2 > 0$.

Proof. We prove the result for Q_2 , the proof for Q_1 is similar. We have

$$\begin{aligned} Q_2(x) \geq & c_0 \int_{\Omega(2x)} \frac{y_1 y_2}{|y|^4} \omega(y) dy + c_0 \int_{\Omega(2x)} \left[G_2^2(x, y) - \frac{y_1 y_2}{|y|^4} \right] \omega(y) dy \\ & + c_0 \int_{[0,1]^2 \setminus \Omega(2x)} G_2^2(x, y) \omega(y) dy - C_1 \|\omega\|_\infty, \end{aligned}$$

throwing away the nonnegative contribution from G_2^1 and estimating Q_2^r by $C_1 \|\omega\|_\infty$ for $x \in D$. First, note that straightforward calculations and estimations give

$$\left| G_2^2(x, y) - \frac{y_1 y_2}{|y|^4} \right| = \frac{|y|^4 y_2 (y_1 - x_1) - |y - x|^2 |y - \bar{x}|^2 y_1 y_2}{|y|^4 |y - x|^2 |y - \bar{x}|^2}.$$

Using

$$\begin{aligned} |y - x|^2 &= |y|^2 - 2x \cdot y + |x|^2 \\ |y - \bar{x}|^2 &= |y|^2 - 2\bar{x} \cdot y + |x|^2 \end{aligned}$$

the nominator can be estimated by

$$\sum_{j=1}^4 |x|^j |y|^{6-j}.$$

For the denominator, note that $y \in \Omega(2x)$ implies that $|y - x| \geq \frac{1}{2}|y|$, $|y - \bar{x}| \geq \frac{1}{2}|y|$, i.e. the denominator is $\gtrsim |y|^8$. Hence, the integral over $\Omega(2x)$ is bounded in absolute value by

$$\|\omega\|_\infty \sum_{j=1}^4 |x|^j \int_{2 \geq |y| \geq 2|x|} |y|^{-2-j} dy \lesssim 1.$$

For the estimation of the integral with domain of integration $[0, 1]^2 \setminus \Omega(2x)$, we distinguish two cases. The more difficult case is given by the condition $x_2 \leq x_1$, and we split the domain of integration into the three parts $[2x_1, 1] \times [0, 2x_2]$, $[0, 2x_1] \times [2x_1, 1]$ and $[0, 2x_1] \times [0, 2x_1]$. For the integral over $[2x_1, 1] \times [0, 2x_2]$, estimate ω by its L^∞ -norm and in the remaining integral we substitute $y_j = x_j + z_j$.

$$\begin{aligned} & \int_{x_1}^{1-x_1} \int_{-x_2}^{x_2} \frac{z_1(x_2 + z_2)}{(z_1^2 + z_2^2)(z_1^2 + (2x_2 + z_2)^2)} dz \leq \int_0^1 \int_{-x_2}^{x_2} \frac{2z_1 x_2}{(z_1^2 + z_2^2)(z_1^2 + x_2^2)} dz_2 dz_1 \\ & \lesssim \int_0^1 \frac{z_1 x_2}{(z_1^2 + x_2^2)} \frac{1}{z_1} \arctan(x_2/z_1) dz_1 \lesssim \arctan(1/x_2) \lesssim 1. \end{aligned}$$

The same strategy for the integral over $[0, 2x_1] \times [2x_1, 1]$ leads to

$$\int_{-x_1}^{x_1} \int_{2x_1-x_2}^{1-x_2} \frac{|z_1|(x_2 + z_2)}{(z_1^2 + z_2^2)(z_1^2 + (2x_2 + z_2)^2)} dz \leq 2 \int_0^{x_1} \int_{x_1}^1 \frac{z_1(x_2 + z_2)}{(z_1^2 + z_2^2)(z_1^2 + (2x_2 + z_2)^2)} dz.$$

Noting

$$\begin{aligned} \int_0^{x_1} \int_{x_1}^1 \frac{z_1 x_2}{(z_1^2 + z_2^2)(z_1^2 + (2x_2 + z_2)^2)} dz &\leq \int_0^{x_1} \int_{x_1}^1 \frac{z_1 x_2}{(z_1^2 + z_2^2)(z_1^2 + x_2^2)} dz_2 dz_1 \\ &\lesssim \int_0^{x_1} \frac{x_2}{z_1^2 + x_2^2} dz_1 \lesssim \arctan(x_1/x_2) \lesssim 1 \end{aligned}$$

and

$$\int_0^{x_1} \int_{x_1}^1 \frac{z_1 z_2}{(z_1^2 + z_2^2)(z_1^2 + (2x_2 + z_2)^2)} dz \leq \int_0^{x_1} \int_{x_1}^1 \frac{z_1 z_2}{(z_1^2 + z_2^2)^2} dz \lesssim 1$$

we can estimate the integral in question by $C\|\omega\|_\infty$.

To estimate the integral over $[0, 2x_1] \times [0, 2x_1]$ first note that

$$\int_{[0, 2x_1] \times [0, 2x_1]} G_2^2(x, y) \omega(y) dy \geq \int_{[0, x_1] \times [0, 2x_1]} G_2^2(x, y) \omega(y) dy.$$

since $\omega \geq 0$ and $G_2^2(x, y) \geq 0$ if $y_1 \leq x_1$. We will estimate the integral over $[0, x_1] \times [0, 2x_1]$ in absolute value, splitting it again into $[0, x_1] \times [0, x_1]$ and $[0, x_1] \times [x_1, 2x_1]$. First, writing $M = M(x, t)$ and using (18) and (63) we get

$$\begin{aligned} \left| \int_{[0, x_1] \times [0, x_1]} G_2^2(x, y) \omega(y) dy \right| &\leq \int_{[0, x_1] \times [0, x_1]} \frac{M y_2^{1-\alpha}}{|y - x| |y - \bar{x}|} dy \\ &\leq \int_{[0, x_1] \times [0, x_1]} \frac{M |y - \bar{x}|^{1-\alpha}}{|y - x| |y - \bar{x}|} dy \leq \int_{[0, x_1] \times [0, x_1]} M |y - x|^{-1-\alpha} dy \\ &\leq \int_{B(x, r)} M |y - x|^{-1-\alpha} dy \leq M r^{1-\alpha} \end{aligned}$$

where $B(x, r)$ is the smallest ball around x containing $[0, 2x_1] \times [0, 2x_1]$. Clearly $r \lesssim x_1$, so the integral is $\lesssim M x_1^{1-\alpha}$.

Next, for the remaining part over $[0, x_1] \times [x_1, 2x_1]$, we estimate ω by $\|\omega\|_\infty$. We need to bound

$$\begin{aligned} \int_{[0, x_1] \times [x_1, 2x_1]} |G_2^2(x, y)| dy &\leq \int_{-x_1}^0 \int_{x_1-x_2}^{2x_1-x_2} \frac{|z_1| z_2}{(z_1^2 + z_2^2)(z_1^2 + (2x_2 + z_2)^2)} dz \\ &\quad + \int_{-x_1}^0 \int_{x_1-x_2}^{2x_1-x_2} \frac{|z_1| x_2}{(z_1^2 + z_2^2)(z_1^2 + (2x_2 + z_2)^2)} dz. \end{aligned}$$

For the integral containing $|z_1| z_2$ we distinguish two cases. In case $x_2 \leq \frac{1}{2}x_1$, we use $z_1^2 + (2x_2 + z_2)^2 \geq z_1^2 + z_2^2$, leading to a bound on the form $\log(1 + \frac{x_1}{x_1-x_2}) \leq C$. If $x_2 \geq \frac{1}{2}x_1$, we use $z_1^2 + (2x_2 + z_2)^2 \geq (x_2 + z_2)^2$ in the denominator and $z_2 \leq (z_2 + x_2)$ in the nominator and get the bound $C x_2^{-1} x_1 \leq C$. The integral with $|z_1| x_2$ is estimated as before.

If $x_1 \leq x_2$, we split $[0, 1]^2 \setminus \Omega(2x)$ into $[0, 1] \times [0, x_2]$, $[0, 2x_1] \times [2x_2, 1]$ and perform similar calculations. In this case, we do not need to use $M(x, t)$. $\square$

Proof of theorem 4.2. Following [8, 11] we observe that the integral

$$\int_{\Omega(2x)} y_1 y_2 |y|^{-4} \omega(y, t) dy$$

can be bounded away from zero by an expression of the form $C_1 \|\omega\|_\infty |\log(1 - \mathbf{m})|$, for $|x| \leq K(1 - \mathbf{m})$. with universal $C_1, K > 0$. Hence we obtain (19). $\square$

4.2. Upper bounds. The following Lemma gives an upper bound on Q_1, Q_2 , in terms of $M_{\widehat{D}}(t)$. Recall that $d(x)$ is the distance to the top of the box, so the upper bound given blows up close to the top of the box. This is, however not a problem, since we mostly have to integrate Q_1, Q_2 along particle trajectories (see the proof Theorem 6.3).

Lemma 4.4. *For $x \in D$,*

$$Q_i(x, t) \lesssim C \|\omega\|_\infty (1 + |\log d(x)|) + M_{\widehat{D}}(t) (|\delta| + \delta_3)^{1-\alpha} \quad (i = 1, 2)$$

Proof. We bound Q_2 , the calculation for Q_1 is analogous. First we note

$$|G_2^k| \lesssim |y - x|^{-1} |y - \bar{x}|^{-1} \quad (k = 1, 2)$$

for $y, x \in [0, 1]^2$. We write $M = M_{\widehat{D}}(t)$, and split the integral in the definition of Q_2 into two parts:

$$\int_{[0,1]^2} G_2^k(x, y) \omega(y) dy = \int_{\widehat{D}} \dots + \int_{[0,1]^2 \setminus \widehat{D}} \dots$$

Since $|\omega(y)| \lesssim M y_2^{1-\alpha}$ and $y_2 \leq |y - \bar{x}|$,

$$\begin{aligned} \left| \int_{\widehat{D}} G_2^k(x, y) \omega(y) dy \right| &\lesssim M \int_{\widehat{D}} y_2^{1-\alpha} |y - x|^{-1} |y - \bar{x}|^{-1} dy \\ &\lesssim M \int_{\widehat{D}} |y - x|^{-1} |y - \bar{x}|^{-\alpha} dy \lesssim M \int_{\widehat{D}} |y - x|^{-1-\alpha} dy \\ &\leq M \int_{B(x, r)} |y - x|^{-1-\alpha} dy \lesssim M r^{1-\alpha} \end{aligned}$$

where $B(x, r)$ is the smallest ball centered at x containing $\widehat{D}$. Obviously $r \lesssim |\delta| + \delta_3$, so the part over $\widehat{D}$ is dominated by $M(|\delta| + \delta_3)^{1-\alpha}$.

For the part over $[0, 1]^2 \setminus \widehat{D}$, we have

$$\begin{aligned} \left| \int_{[0,1]^2 \setminus \widehat{D}} G_2^k(x, y) \omega(y) dy \right| &\lesssim \|\omega\|_\infty \int_{[0,1]^2 \setminus \widehat{D}} |y - x|^{-2} dy \\ &\lesssim \|\omega\|_\infty \int_{B(x, 10) \setminus B(x, d(x))} |y - x|^{-2} dy \lesssim \|\omega\|_\infty |\log d(x)| \end{aligned}$$

where we have used $|G_2^k| \lesssim |x - y|^{-1} |y - \bar{x}|^{-1}$ and $|y - \bar{x}| \geq |y - x|$ for $x, y \in [0, 1]^2$. Note also that for $x \in D$, $[0, 1]^2 \setminus \widehat{D}$ is completely contained in $B(x, 10) \setminus B(x, d(x))$ because of (17).

For Q_i^r we have the estimate $|Q_i^r(x, t)| \leq C \|\omega\|_\infty$, concluding the proof. $\square$

The following important Lemma allows us to control the coefficients of the ODE system (8) in terms of the quantity $M_{\widehat{D}}$. Recall that $d(x)$ is the distance from $x \in D$ to the top of the box.

Lemma 4.5. *We have the following estimates for $x \in D$:*

$$\begin{aligned} |c(x, t)| &\leq C(\alpha)M_{\widehat{D}}(t)x_2^{1-\alpha} + C(\alpha, \gamma_1, \gamma_2)x_2^{1-\gamma_1-\gamma_2}x_1^{\gamma_2}d(x)^{-1+\gamma_1}, \\ |b(x, t)| &\leq C(\alpha)M_{\widehat{D}}(t)x_1^{1-\alpha}(1 + |\log d(x)|) + C(\alpha, \gamma)x_1^{1-\gamma}d(x)^{-1+\gamma}, \\ \left| x_i \frac{\partial Q_i}{\partial x_i}(x, t) \right| &\leq C(\alpha)M_{\widehat{D}}(t)x_i^{1-\alpha}(1 + |\log d(x)|) + C(\alpha, \gamma)x_i^{1-\gamma}d(x)^{-1+\gamma} \end{aligned}$$

where $\gamma, \gamma_1, \gamma_2 \in (0, 1)$, $\gamma_1 + \gamma_2 < 1$, $i = 1, 2$ and the constants do not depend on $\delta_1, \delta_2, \delta_3, t$.

Proof. This is a consequence of Proposition 9.9 (see appendix) and the definition of $c, b, x_i \partial_{x_i} Q_i$ (see (15)). Note that we have

$$\left| \frac{\partial Q_i^r}{\partial x_j}(x, t) \right| \leq C\|\omega\|_\infty$$

for $x \in D$. When we estimate e.g. c , we encounter a term of the form $x_2 \frac{\partial Q_2^r}{\partial x_1}(x, t)$, obtaining a bound of the form $Cx_2\|\omega\|_\infty$, which can be absorbed into

$$C(\alpha)M_{\widehat{D}}(t)x_2^{1-\alpha}.$$

□

Remark 4.6. It is not possible to set $\alpha = 0$ in the estimates of Lemma 4.5, i.e. if we replace $M_{\widehat{D}}$ by $\|\nabla\omega\|_{D,\infty}$, then e.g. the first term on the right-hand side of the estimate for c would contain a logarithmic expression

$$\|\nabla\omega\|_{D,\infty}x_2|\log x_2|.$$

This is the main reason why we do not adopt the stronger feeding condition (10), since our main argument cannot be applied to this kind of logarithmic terms.

5. PERTURBATION THEORY FOR A SYSTEM OF ORDINARY DIFFERENTIAL EQUATIONS

In this section we derive estimates for an ODE system of the form

$$\dot{\xi}(t) = \begin{pmatrix} a(t) & c(t) \\ b(t) & -a(t) \end{pmatrix} \xi(t)$$

where a, b, c are given smooth functions on a time interval $[T_0, T_e]$. This part is independent of the actual structure of a, b, c from the ODE (8).

The idea will be to perturb from the system with $c \equiv 0$, which can be solved explicitly. We write

$$P(t) := \begin{pmatrix} a(t) & 0 \\ b(t) & -a(t) \end{pmatrix}, \quad S(t) := \begin{pmatrix} 0 & c(t) \\ 0 & 0 \end{pmatrix}.$$

Definition 5.1. Let the integral operators $\widehat{P}, \widehat{S}$ be given by

$$(\widehat{P}\xi)(t) = \int_{T_0}^t P(s)\xi(s) ds, \quad (\widehat{S}\xi)(t) = \int_{T_0}^t S(s)\xi(s) ds.$$

Recall that $A(t) = \int_{T_0}^t a(s) ds$. It is convenient to introduce the following operators:

$$\begin{aligned} (F^+g)(t) &= g(t) + e^{A(t)} \int_{T_0}^t a(s) e^{-A(s)} g(s) ds, \\ (F^-g)(t) &= g(t) - e^{-A(t)} \int_{T_0}^t a(s) e^{A(s)} g(s) ds. \end{aligned}$$

Proposition 5.2. (a) *The operator $(I - \widehat{P})$ is bounded and bijective as an operator from $C[T_0, T]$ into $C[T_0, T]$.*

(b) *Consider the Volterra integral equation*

$$(20) \quad \phi = \widehat{P}\phi + g$$

with given $g \in C([T_0, T], \mathbb{R}^2)$. The solution $\phi = (I - \widehat{P})^{-1}g$ is given by

$$(21) \quad \begin{aligned} \phi_1(t) &= F^+g_1 \\ \phi_2(t) &= F^-g_2 + e^{-A} \int_{T_0}^t e^A b F^+g_1 ds \end{aligned}$$

Proof. The statement (a) is standard. Statement (b) is an easy calculation, noting that (20) is equivalent to the ODE system $\dot{\xi} = P\xi + \dot{g}$ for $g \in C^1$. $\square$

The initial value problem for the system

$$\dot{\xi} = (P + S)\xi, \quad \xi(T_0) \text{ given}$$

is equivalent to the Volterra integral equation

$$(22) \quad \xi = (\widehat{P} + \widehat{S})\xi + \xi(T_0).$$

We can write $\xi = (I - \widehat{P})^{-1}\mathbf{w}$ for some $\mathbf{w} \in C[T_0, T]$. This leads to

$$(23) \quad \mathbf{w} = \widehat{S}(I - \widehat{P})^{-1}\mathbf{w} + \xi(T_0).$$

The following proposition gives a representation of the solution ξ in terms of $\mathbf{w}$:

Proposition 5.3. *Let $\xi \in C[T_0, T]$ solve the integral equation (22) with given $\xi(T_0)$. Then*

$$(24) \quad \begin{aligned} \xi_1(t) &= (F^+w_1)(t), \quad \xi_2(t) = \xi_2(T_0)e^{-A} + e^{-A} \int_{T_0}^t e^A b F^+w_1 ds \\ w_1(t) &= \xi_1(T_0) + \xi_2(T_0) \int_{T_0}^t e^{-A} c ds + \int_{T_0}^t e^{-A} c \int_{T_0}^s e^A b F^+w_1 d\tau ds, \\ w_2(t) &= \xi_2(T_0) \end{aligned}$$

Proof. First note that

$$(25) \quad \widehat{S}(I - \widehat{P})^{-1}\mathbf{w} = \widehat{S}\xi = \left(\int_{T_0}^t c(s)\xi_2(s) ds, 0 \right)$$

and hence by (23), $w_2(t) = \xi_2(T_0)$. It is easy to compute F^-w_2 :

$$(26) \quad F^-w_2 = F^-\xi_2(T_0) = \xi_2(T_0)e^{-A}.$$

Recalling $\boldsymbol{\xi} = (I - \widehat{P})^{-1} \mathbf{w}$ we get from (20) and (21) with $g = \mathbf{w}$ and $\phi = \boldsymbol{\xi}$ and using (26)

$$(27) \quad \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} = \boldsymbol{\xi} = (I - \widehat{P})^{-1} \mathbf{w} = \begin{pmatrix} F^+ w_1 \\ \xi_2(T_0) e^{-A} + e^{-A} \int_{T_0}^t e^A b F^+ w_1 ds \end{pmatrix}$$

which is the first line of (24). From (25) follows

$$\begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = \widehat{S} \boldsymbol{\xi} + \boldsymbol{\xi}(T_0) = \begin{pmatrix} \int_{T_0}^t c \xi_2 ds + \xi_1(T_0) \\ \xi_2(T_0) \end{pmatrix}.$$

Together with (27) we get

$$\begin{aligned} w_1 &= \int_{T_0}^t c \xi_2 ds + \xi_1(T_0) = \int_{T_0}^t c \left(\xi_2(T_0) e^{-A} + e^{-A} \int_{T_0}^s e^A b F^+ w_1 d\tau \right) ds + \xi_1(T_0) \\ &= \xi_1(T_0) + \xi_2(T_0) \int_{T_0}^t e^{-A} c ds + \int_{T_0}^t e^{-A} c \int_{T_0}^s e^A b F^+ w_1 d\tau ds \end{aligned}$$

□

We will need the following Gronwall-type inequality by Wilett [10, 9]:

Lemma 5.4. *Let $z, f_0, f_1, f_2, v_1, v_2$ are nonnegative, integrable functions on $[T_0, T]$ and suppose z satisfies the following integral inequality:*

$$z(t) \leq f_0(t) + f_1(t) \int_{T_0}^t v_1(s) z(s) ds + f_2(t) \int_{T_0}^t v_2(s) z(s) ds.$$

Then $z \leq H f_0$, where H is the following functional

$$\begin{aligned} (H f_0)(t) &= f_0 + f_1 \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 f_0 \\ &\quad + \left[f_2(t) + f_1(t) \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 f_2 \right] \\ (28) \quad &\quad \times \exp \left(\int_{T_0}^t v_2 \left[f_2(s) + f_1(s) \exp \left(\int_{T_0}^s v_1 f_1 \right) \int_{T_0}^s v_1 f_2 \right] \right) \\ &\quad \times \int_{T_0}^t v_2 \left[f_0(s) + f_1(s) \exp \left(\int_{T_0}^s v_1 f_1 \right) \int_{T_0}^s v_1 f_0 \right] \end{aligned}$$

We write $H f_0$ to emphasize the linear dependency on f_0 .

Proof. We give the proof for reference. Recall first the following basic form of Gronwall's integral inequality [9]. Suppose z, r, f_1, v_1 are nonnegative functions on $[T_0, T]$ satisfying the integral inequality

$$z(t) \leq r(t) + f_1(t) \int_{T_0}^t v_1 z ds,$$

then

$$(29) \quad z(t) \leq r(t) + f_1(t) \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 r ds \quad (t \in [T_0, T]).$$

Set $r = f_0 + f_2 \int_{T_0}^t v_2 z$ and apply (29). This leads to the following bound for z :

$$(30) \quad z(t) \leq f_0 + f_2 \int_{T_0}^t v_2 z + f_1(t) \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 \left[f_0 + f_2 \int_{T_0}^s v_2 z \right].$$

Note that

$$\int_{T_0}^t v_1 f_2 \int_{T_0}^s v_2 z \leq \left(\int_{T_0}^t v_1 f_2 \right) \int_{T_0}^t v_2 z$$

since $v_1, f_2, z, v_2 \geq 0$. Thus (30) implies

$$\begin{aligned} z(t) &\leq f_0 + f_1 \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 f_0 + \\ &\quad + \left[f_2(t) + f_1(t) \exp \left(\int_{T_0}^t v_1 f_1 \right) \left(\int_{T_0}^t v_1 f_2 \right) \right] \int_{T_0}^t v_2 z. \end{aligned}$$

Applying (29) again, this time with $r = f_0 + f_1 \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 f_0$, yields the result (28). $\square$

Lemma 5.5. *Let $\xi \in C[T_0, T]$ solve the integral equation (22) with given $\xi(T_0)$. Then the estimates*

$$(31) \quad \begin{aligned} |\xi_1(t)| &\leq (Hf_0)(t) + e^A \int_{T_0}^t v_1 Hf_0 \\ |\xi_2(t)| &\leq |e^{-A}\xi_2(T_0)| + e^{-A} \left[\int_{T_0}^t v_2 Hf_0 + \int_{T_0}^t e^{2A}|b| \int_{T_0}^s v_1 Hf_0 \right], \end{aligned}$$

hold, where H is the functional (28) and where

$$\begin{aligned} f_1(t) &= \int_{T_0}^t e^{-A}|c| \int_{T_0}^s e^{2A}|b|, & f_2(t) &= \int_{T_0}^t e^{-A}|c|, \\ f_0(t) &= |\xi_1(T_0)| + f_2(t)|\xi_2(T_0)|, & v_1(t) &= |a(t)|e^{-A}, \\ v_2(t) &= |b(t)|e^A. \end{aligned}$$

Proof. Using obvious estimations, we get from (24) the following integral inequality for $|w_1|$:

$$\begin{aligned} |w_1(t)| &\leq |\xi_1(T_0)| + |\xi_2(T_0)| \int_{T_0}^t e^{-A}|c| + \int_{T_0}^t e^{-A}|c|ds \int_{T_0}^t e^A|b||w_1|ds \\ &\quad + \int_{T_0}^t e^{-A}|c| \int_{T_0}^s e^{2A}|b|d\tau ds \int_{T_0}^t |a|e^{-A}|w_1| \\ &= f_0(t) + f_1(t) \int_{T_0}^t v_1 |w_1|ds + f_2(t) \int_{T_0}^t v_2 |w_1|ds, \end{aligned}$$

where the expressions f_0, f_1, f_2, v_1, v_2 are given as in the statement of the Lemma. Now using Lemma 5.4, we obtain $|w_1(t)| \leq Hf_0$ on $[T_0, T]$. The inequalities (31) follow from the formulas (21). $\square$

Remark 5.6. The reader might wonder why we did not perturb from a diagonal system, i.e. regard also b as a perturbation like c as in the heuristic discussion. While it is certainly possible to derive corresponding perturbation formulas for ξ_1, ξ_2 , it turns out that the balance of growing and decaying factors is not favorable for the arguments in Section 6. Fortunately, the perturbation from the tridiagonal system behaves in a more stable way.

6. MAIN ARGUMENT

6.1. The main technical result. In order to formulate our main technical result, we introduce a notion of *harmless nonlinear bound*.

Definition 6.1. A function $\mathcal{N} = \mathcal{N}(R, \beta, \alpha, \boldsymbol{\delta}, M)$ where all arguments are nonnegative numbers is a *harmless nonlinear function* if for fixed $\alpha \in (0, 1), \beta > 0$ the following holds: For any given $R > 0$, there exists $\bar{\delta}_2(R) > 0$ and a number $\bar{\delta}_1 = \bar{\delta}_1(R, \bar{\delta}_2) > 0$ such that for all $\delta_2 \leq \bar{\delta}_2, \delta_1 \leq \bar{\delta}_1$ the inequality

$$\mathcal{N}(R, \beta, \alpha, \boldsymbol{\delta}, R) < R$$

holds.

Recall the box $\widehat{D}$ is said to satisfy the conditions of *controlled feeding* if there is a $R \geq 0$ with

$$|\partial_{x_1}\omega(x, t)| \leq Rx_2^{1-\alpha}, \quad |\partial_{x_2}\omega(x, t)| \leq R \quad (x \in \widehat{D} \setminus D)$$

for all times $t \geq 0$. R is called *feeding parameter*. For convenience, we introduce the following definition.

Definition 6.2. Let $T > 0, \beta > 0$. We say that the flow is β -hyperbolic in the box D on $[0, T]$ if

$$Q_i(x, t) \geq \beta \quad (x \in D, t \in [0, T], i = 1, 2).$$

Theorem 6.3. Let $0 < \alpha < 1/4$. There exists a harmless nonlinear function $\mathcal{N} = \mathcal{N}(R, \beta, \alpha, \boldsymbol{\delta}, M)$ (determined by a-priori known data) with the following properties. If ω is a solution of the Euler equation, $\widehat{D}$ a box defined by (16) with parameters $\delta_1, \delta_2, \delta_3 > 0$ satisfying (17) and $T > 0$ is such that

- (i) the flow is β -hyperbolic in the box D on the time interval $[0, T]$,
- (ii) the box $\widehat{D}$ satisfies the conditions of controlled feeding with parameter $R > \|\omega\|_\infty$,
- (iii) the initial data satisfies,

$$M_D(0) < R, \quad \left| \frac{\partial \omega_0}{\partial x_1}(x) \right| \leq Rx_2^{1-\alpha}, \quad \left| \frac{\partial \omega_0}{\partial x_2}(x) \right| \leq R \quad (x \in D),$$

- (iv) there exists a number K such that

$$M_D(t) \leq K \quad (t \in [0, T]),$$

then

$$M_D(t) \leq \mathcal{N}(R, \alpha, \beta, \boldsymbol{\delta}, K) \quad (t \in [0, T])$$

holds.

6.2. Estimates along particle trajectories. In this section we develop the technical tools to prove Theorem 6.3. The proofs for the estimates for f_0, f_1, f_2 and Hf_0 along trajectories are heavily interconnected (see Figure 3). We advise the reader to concentrate on the main flow of arguments indicated by the bold arrows and boxes in the map of Section 6.




FIGURE 3. Map of Section 6.

Let ω be a given double-odd solution of the Euler equation that is in $C^1([0, \infty), C^2(\mathbb{T}))$. Moreover, let $\widehat{D}$ be a box depending on the parameters $\delta_1, \delta_2, \delta_3 > 0$ satisfying the conditions (17).

Suppose also that for the remainder of this section, (i)-(iv) from theorem 6.3 are satisfied. For abbreviation, we write in the following

$$M := \max\{K, R\}.$$

We observe the following important fact: since $\delta_1, \delta_2, \delta_3 \leq 1$,

$$(32) \quad M_{\widehat{D}}(t) \leq M$$

holds.

We consider associated particle trajectories, which are the solutions of

$$(33) \quad \dot{X}_1 = -X_1 Q_1, \quad \dot{X}_2 = X_2 Q_2.$$

More precisely, we define the particle trajectories as follows: for any $(x_0, t_0) \in \overline{D} \times [0, \infty)$ we take the maximal solution of $t \mapsto \mathbf{X}(t)$ of (33) which passes through (x_0, t_0) , and lies $\overline{D}$. $\mathbf{X}$ is defined on an interval $[T_0, T_e]$ such that

- (i) $\mathbf{X}(t) \in \overline{D}$ for all $T_0 \leq t \leq T_e$,
- (ii) either $T_0 = 0$ or $T_0 > 0$, in which case necessarily $\mathbf{X}(T_0) \in \partial D$,
- (iii) $\mathbf{X}(T_e) \in \partial D$.

Observe that $\mathbf{X}$ is given by

$$(34) \quad \begin{aligned} X_1(t) &= X_1(T_0) \exp \left(- \int_{T_0}^t Q_1(\mathbf{X}(s), s) ds \right) \\ X_2(t) &= X_2(T_0) \exp \left(\int_{T_0}^t Q_2(\mathbf{X}(s), s) ds \right). \end{aligned}$$

We call T_0 the entry time and T_e the exit time of a particle trajectory. $T_0 = 0$ if the particle starts in $\overline{D}$ for $t = 0$.

The next proposition gives a upper bound for the time a particle can spend in the upper half of the box D , provided the flow is β -hyperbolic.

Proposition 6.4. *Suppose that the flow is β -hyperbolic in the box D on the time interval $[0, T]$. Let $\mathbf{X}$ be a particle trajectory whose entry time T_0 is smaller than T . Then if $X_2(T_0) \neq 0$ there is either a time $T_1, T_e > T_1 \geq T_0$ such that*

$$X_2(t) \geq \frac{1}{2} \delta_2 \quad (t \in [T_1, T])$$

or

$$X_2(t) \leq \frac{1}{2} \delta_2 \quad (t \in [T_0, T]).$$

If T_1 exists, we have the estimate

$$T_e - T_1 \leq \beta^{-1} \log(2).$$

Proof. The statement on the time T_1 follows directly from the fact that the flow is β -hyperbolic in the box. If T_1 exists, we have analogously to (34)

$$\delta_2 = X_2(T_e) = X_2(T_1) \exp \left(\int_{T_1}^{T_e} Q_2 ds \right) \geq \frac{\delta_2}{2} \exp \left(\int_{T_1}^{T_e} Q_2 ds \right) \geq \frac{\delta_2}{2} \exp(\beta(T_e - T_1)).$$

Solving for $T_e - T_1$ gives the result. $\square$

Definition 6.5. We call a function $g = g(\alpha, \beta, \delta, M)$ *harmless generic factor* it has the following property: There exists a $p_0 > 0$ such that for all $p \geq p_0$ and fixed α, β, M

$$g(\alpha, \beta, \delta_2^p, \delta_2, M)$$

is bounded as $\delta_2 \rightarrow 0$.

Remark 6.6. For example, a function of the form

$$g = C(\alpha, \beta) [\delta_2^{\gamma_3} M(1 + |\log \delta_2|) + \delta_1^{\gamma_1} \delta_2^{-\gamma_2} + 1]^{\gamma_4} + C(\alpha, \beta, \gamma_j)$$

($\gamma_j > 0$) is a harmless generic factor, and e^g is also a harmless generic factor if g is one. When performing estimations, we shall often absorb harmless generic factors into one another, so the actual meaning of g may change from line to line.

In our argument there will appear only finitely many different generic factors (although all denoted by g). To make the boundedness of them all work as $\delta_2 \rightarrow 0$ we just pick a p that is bigger than all the p_0 of all appearing generic factors.

Our goal will be to obtain estimates for the quantities f_0, f_1, f_2, v_1, v_2 along a single particle trajectory, up to the given time T , so that we can apply our ODE estimates from section 5. The crucial point is that our bounds depend *not directly* on ω , T , T_e but only on $\beta, \alpha, \mathbf{X}(T_0)$. For the estimations below we often refer to a fixed particle trajectory with entry time T_0 , along which we evaluate integrals over time of the quantities Q_1, Q_2, c etc. To make the notation more compact, we often skip $\mathbf{X}$ in the arguments of the integrands, e.g. we write

$$\int_{T_0}^t |c| e^{-A} ds = \int_{T_0}^t |c| e^{-A(s)} ds = \int_{T_0}^t |c(\mathbf{X}(s), s)| \exp \left(\int_{T_0}^s a(\mathbf{X}(\tau), \tau) d\tau \right) ds.$$

Lemma 6.7. *For any $t \leq T_e$,*

$$X_2(T_0) \leq \delta_2 \exp \left(- \int_{T_0}^t Q_2(\mathbf{X}(s), s) ds \right).$$

Proof. Since the particle trajectory lies in D for $t \in [T_0, T_e]$,

$$\delta_2 \geq X_2(t) = X_2(T_0) \exp \left(\int_{T_0}^t Q_2 ds \right)$$

holds. □

Let $\phi : [0, \infty) \rightarrow [0, \infty)$ be a function with the properties

$$\phi(s) \leq 1 - e^{-s}$$

and ϕ monotone nondecreasing on $[0, \infty)$, ϕ linear on $[0, s^*]$ and ϕ constant on $[s^*, \infty)$ for some s^* . We fix such a function ϕ for the following.

Proposition 6.8. *Along a particle trajectory in a β -hyperbolic flow in D , we have the following for $t \in [T_0, \min\{T_e, T\}]$:*

(i)

$$\begin{aligned} X_1(t) &\leq \delta_1 \exp(-\beta(t - T_0)) \\ X_2(t) &\leq \delta_2 \exp(-\beta(\min\{T_e, T\} - t)), \end{aligned}$$

(ii)

$$d(\mathbf{X}(t)) \geq \delta_2 \phi \left(\int_t^{\min\{T_e, T\}} Q_2 ds \right) \geq \delta_2 \phi(\beta(\min\{T, T_e\} - t)),$$

(iii) *Suppose T_1 from proposition 6.4 exists. Then the following holds for any $\gamma \in (0, 1)$ and $t \in [T_1, \min\{T_e, T\}]$:*

$$\begin{aligned} \int_{T_1}^t d(\mathbf{X}(s))^{-1+\gamma} ds &\leq C(\gamma, \beta) \delta_2^{-1+\gamma}, \\ \int_{T_1}^t |\log d(\mathbf{X}(s))| ds &\leq C(\beta) |\log \delta_2| \end{aligned}$$

with $C(\beta), C(\gamma, \beta)$ independent of the trajectory.

Proof. For (i), recall that under the assumption of β -hyperbolic flow, $Q_2 \geq \beta$. From (34), we get

$$\begin{aligned}
 X_2(t) &= X_2(T_0) \exp \left(\int_{T_0}^{\min\{T_e, T\}} Q_2 \, ds - \int_t^{\min\{T_e, T\}} Q_2 \, ds \right) \\
 &= X_2(\min\{T_e, T\}) \exp \left(- \int_t^{\min\{T_e, T\}} Q_2 \, ds \right) \\
 (35) \quad &\leq \delta_2 \exp(-\beta(\min\{T_e, T\} - t)),
 \end{aligned}$$

noting that $X_2(\min\{T_e, T\}) \leq \delta_2$. The bound for X_1 is analogous.

Now we show (ii). Recall that $d(\mathbf{X}) = \delta_2 - X_2(t)$. Hence by (35)

$$\begin{aligned}
 \delta_2 - X_2(t) &= \delta_2 \left(1 - \exp \left(- \int_t^{\min\{T_e, T\}} Q_2 \, ds \right) \right) \geq \delta_2 \phi \left(\int_t^{\min\{T_e, T\}} Q_2 \, ds \right) \\
 &\geq \delta_2 \phi(\beta(\min\{T_e, T\} - t)).
 \end{aligned}$$

(iii) We split the integrals by introducing the time T^* defined as follows: T^* is the maximum of all $s \in [T_1, \min\{T_e, T\}]$ such that

$$\phi(\beta(\min\{T_e, T\} - s)) = \phi(s^*).$$

If there are no such s , we set $T^* = T_1$. Thus we split the integrals in (iii) as follows:

$$\int_{T_1}^t = \int_{T_1}^{T^*} \dots + \int_{T^*}^t \dots$$

if $t \geq T^*$, otherwise we have only one integral from T_1 to t . We calculate

$$\begin{aligned}
 \int_{T_1}^{T^*} d(\mathbf{X}(s))^{-1+\gamma} \, ds &\leq \delta_2^{-1+\gamma} \int_{T_1}^{T^*} \phi(\beta(\min\{T_e, T\} - s))^{-1+\gamma} \, ds \\
 &\leq \delta_2^{-1+\gamma} (T_e - T_1) \phi(s^*)^{-1+\gamma} \leq C(\beta, \gamma) \delta_2^{-1+\gamma} \\
 \int_{T^*}^t d(\mathbf{X}(s))^{-1+\gamma} \, ds &\leq \delta_2^{-1+\gamma} \int_{T^*}^t \phi(\beta(\min\{T_e, T\} - s))^{-1+\gamma} \, ds \\
 &\lesssim \delta_2^{-1+\gamma} \beta^{-1+\gamma} \int_{T^*}^t (\min\{T_e, T\} - s)^{-1+\gamma} \, ds \\
 &\lesssim \delta_2^{-1+\gamma} \beta^{-1+\gamma} \int_{T_1}^{\min\{T_e, T\}} (\min\{T_e, T\} - s)^{-1+\gamma} \, ds \\
 &\lesssim \delta_2^{-1+\gamma} \beta^{-1+\gamma} \int_0^{T_e - T_1} z^{-1+\gamma} \, dz \lesssim \delta_2^{-1+\gamma} C(\beta, \gamma).
 \end{aligned}$$

using (ii), Proposition 6.4 to estimate $T_e - T_1$ and the fact that ϕ is linear on $[0, s^*]$. The second integral is treated analogously. $\square$

Lemma 6.9. *Along a particle trajectory, we have, for $T_0 \leq t \leq \min\{T, T_e\}$,*

$$e^{\pm A(t)} \leq g(\alpha, \beta, \boldsymbol{\delta}, M) \exp \left(\pm \int_{T_0}^t Q_2(s) ds \right),$$

$$\exp \left(\pm \int_{T_0}^t Q_i(s) ds \right) \leq g(\alpha, \beta, \boldsymbol{\delta}, M) \exp \left(\pm \int_{T_0}^t Q_j(s) ds \right), \quad i, j = 1, 2$$

where $g(\alpha, \beta, \boldsymbol{\delta}, M)$ are harmless generic factors depending only on the quantities indicated.

Proof. We prove the first inequality of the Lemma, for the other we use similar arguments. Recall $a(t) = Q_2(t) + X_2(t)\partial_{x_2}Q_2(t)$, $A(t) = \int_{T_0}^t a(s) ds$ and thus

$$\pm A(t) \leq \pm \int_{T_0}^t Q_2(s) ds + \int_{T_0}^t |X_2(s)\partial_{x_2}Q_2(s)| ds.$$

We now use Lemma 4.5 and (32):

$$\begin{aligned} \int_{T_0}^t |X_2(s)\partial_{x_2}Q_2(s)| ds &\leq C(\alpha)M \int_{T_0}^{\min\{T, T_e\}} X_2^{1-\alpha}(1 + |\log d(\mathbf{X})|) ds \\ &\quad + C(\alpha, \gamma) \int_{T_0}^{\min\{T, T_e\}} X_2^{1-\gamma} d(\mathbf{X})^{-1+\gamma} ds. \end{aligned}$$

Note that the interval of integration has been enlarged. With T_1 from proposition 6.4 we split the interval of integration into $[T_0, T_1]$ and $[T_1, \min\{T, T_e\}]$ provided $\min\{T, T_e\} \geq T_1$. The case $\min\{T, T_e\} < T_1$ is analogous.

In the part over $[T_0, T_1]$, while $d(\mathbf{X}) \geq \frac{1}{2}\delta_2$, we cannot control the length of the time interval, so we estimate as follows:

$$\begin{aligned} \int_{T_0}^{T_1} X_2^{1-\alpha}(1 + |\log d(\mathbf{X})|) &\leq \delta_2^{1-\alpha} \int_{T_0}^{T_1} e^{-(1-\alpha)\beta(\min\{T, T_e\}-s)} (C + |\log \delta_2|) ds \\ &\leq C\delta_2^{1-\alpha} |\log \delta_2| \int_0^\infty e^{-(1-\alpha)\beta z} dz \\ &\leq C(\alpha, \beta)\delta_2^{1-\alpha} |\log \delta_2|, \end{aligned}$$

using part (i) of proposition 6.8 and $d(\mathbf{X}(s)) \geq \frac{1}{2}\delta_2$ for $s \in [T_0, T_1]$, and δ_2 sufficiently small. In the part over $[T_1, \min\{T, T_e\}]$ the length of the time interval is bounded but $|\log d(\mathbf{X})|$ is unbounded, so we proceed differently:

$$\begin{aligned} \int_{T_1}^{\min\{T, T_e\}} X_2^{1-\alpha}(1 + |\log d(\mathbf{X})|) &\leq \delta_2^{1-\alpha} \int_{T_1}^{\min\{T, T_e\}} |\log d(\mathbf{X})| ds \\ &\leq C(\beta)\delta_2^{1-\alpha} |\log \delta_2|. \end{aligned}$$

using statement (iii) of Proposition 6.8 and $X_2 \leq \delta_2$.

For the second integral involving $X_2^{1-\gamma} d(\mathbf{X})^{-1+\gamma}$, we note

$$\begin{aligned} \int_{T_0}^{T_1} X_2^{1-\gamma} d(\mathbf{X})^{-1+\gamma} &\leq C(\gamma) \delta_2^{-1+\gamma} \int_{T_0}^{T_1} (\delta_2 e^{-\beta(\min\{T, T_e\}-s)})^{1-\gamma} ds \\ &\leq C(\gamma, \beta) \\ \int_{T_1}^{\min\{T, T_e\}} X_2^{1-\gamma} d(\mathbf{X})^{-1+\gamma} &\leq \delta_2^{1-\gamma} \int_{T_1}^{\min\{T, T_e\}} d(\mathbf{X})^{-1+\gamma} \\ &\leq C(\gamma, \beta) \end{aligned}$$

by proposition 6.8, (i) and (iii) and moreover using $X_2 \leq \delta_2$. This yields finally

$$\int_{T_0}^t |X_2(s) \partial_{x_2} Q_2(s)| ds \leq [C(\alpha, \beta) M \delta_2^{1-\alpha} |\log \delta_2| + C(\gamma, \beta)]$$

implying the result, since the term in square brackets is a harmless generic factor.

To prove the second inequality, we use (the velocity field is divergence-free)

$$Q_1(t) + X_1(t) \partial_{x_1} Q_1(t) = Q_2(t) + X_2(t) \partial_{x_2} Q_2(t)$$

implying $|Q_i| \leq |Q_j| + \sum_{k=1}^2 |x_k \partial_{x_k} Q_k|$. The expressions involving $x_k \partial_{x_k} Q_k$ are estimated as before. $\square$

6.3. Estimates for f_0, f_1, f_2, v_1, v_2 and Hf_0 .

Lemma 6.10. *The following estimates hold for $T_0 \leq t \leq \min\{T, T_e\}$:*

$$\begin{aligned} (36) \quad f_2(t) &\leq g(\alpha, \beta, \boldsymbol{\delta}, M) X_2(T_0)^{1-\alpha}, \\ f_0(t) &\leq R g(\alpha, \beta, \boldsymbol{\delta}, M) X_2(T_0)^{1-\alpha}. \end{aligned}$$

Proof. We write $g = g(\alpha, \beta, \boldsymbol{\delta}, M)$ for any occurring harmless factor. Using Lemma 4.5 with $\gamma_1 = \gamma_2 = \frac{\alpha}{2}$,

$$\begin{aligned} f_2(t) &= \int_{T_0}^t e^{-A} |c| \lesssim M \int_{T_0}^{\min\{T, T_e\}} e^{-A} X_2^{1-\alpha} ds \\ &\quad + C(\alpha) \int_{T_0}^{\min\{T, T_e\}} e^{-A} X_2^{1-\alpha} X_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2} ds. \end{aligned}$$

Observe first that by Lemma 6.9 $e^{-A(s)}$ is estimated by $\exp\left(-\int_{T_0}^s Q_2 d\tau\right)$ and thus using $Q_2 \geq \beta$ again we get

$$(37) \quad e^{-A} X_2(s)^{1-\alpha} \leq g X_2(T_0)^{1-\alpha} \exp\left(-\alpha \int_{T_0}^s Q_2 d\tau\right) \leq g X_2(T_0)^{1-\alpha} \exp(-\alpha \beta(s - T_0)).$$

Employing (37) to estimate the integral containing $e^A X_2^{1-\alpha}$ yields:

$$\int_{T_0}^{\min\{T, T_e\}} e^{-A} X_2^{1-\alpha} ds \leq g X_2(T_0)^{1-\alpha} \int_{T_0}^{\infty} e^{-\alpha \beta(s - T_0)} ds \leq g X_2(T_0)^{1-\alpha} C(\alpha, \beta).$$

For the integral containing $e^{-A} X_2^{1-\alpha} X_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2}$, we use (37) again and estimate

$$\int_{T_0}^{\min\{T, T_e\}} e^{-A} X_2^{1-\alpha} X_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2} ds \leq g X_2(T_0)^{1-\alpha} \delta_1^{\alpha/2} \int_{T_0}^{\min\{T, T_e\}} e^{-\alpha \beta(s - T_0)} d(\mathbf{X})^{-1+\alpha/2} ds.$$

As in the proof of Lemma 6.9, we split the interval of integration into $[T_0, T_1]$ and $[T_1, \min\{T, T_e\}]$ in case $T_1 \leq \min\{T, T_e\}$, obtaining

$$(38) \quad \int_{T_0}^{T_1} e^{-\alpha\beta(s-T_0)} d(\mathbf{X})^{-1+\alpha/2} ds \lesssim \delta_2^{-1+\alpha/2},$$

$$(39) \quad \int_{T_1}^{\min\{T, T_e\}} e^{-\alpha\beta(s-T_0)} d(\mathbf{X})^{-1+\alpha/2} ds \lesssim \int_{T_1}^{\min\{T, T_e\}} d(\mathbf{X})^{-1+\alpha/2} ds \lesssim \delta_2^{-1+\alpha/2}$$

where we have used $d(\mathbf{X}) \geq \frac{1}{2}\delta_2$ for (38) and $e^{-\alpha\beta(s-T_0)} \leq 1$ and Proposition 6.8 for (39). The case $T_1 \geq \min\{T, T_e\}$ is covered by (38). To estimate f_0 , we use that the feeding condition holds and that assumption (iii) from Theorem 6.3 holds. This gives

$$\begin{aligned} |\xi_1(T_0)| &= |\partial_{x_1}\omega(\mathbf{X}(T_0), T_0)| \leq R X_2(T_0)^{1-\alpha}, \\ |\xi_2(T_0)| &= |\partial_{x_2}\omega(\mathbf{X}(T_0), T_0)| \leq R \end{aligned}$$

for both of the cases $T_0 = 0$ (particle starts in D) and $T_0 > 0$ (particle starts in or crosses the feeding zone before entering D). Now use the definition of f_0 and the estimate (36) for f_2 . $\square$

Lemma 6.11. *For $T_0 \leq t \leq \min\{T, T_e\}$,*

$$f_1(t) \leq g(\alpha, \beta, \boldsymbol{\delta}, M) \delta_1^{1-\alpha} \delta_2^\alpha e^{\alpha \int_{T_0}^t Q_2 ds}$$

with a harmless generic factor g depending on the quantities indicated.

Proof. We abbreviate again $g = g(\alpha, \beta, \boldsymbol{\delta}, M)$. First we claim that for sufficiently small δ_2

$$(40) \quad \int_{T_0}^t e^{2A} |b| ds \leq g X_1(T_0)^{1-\alpha} \left[M |\log \delta_2| + \delta_1^{\frac{\alpha}{2}} \delta_2^{-1+\frac{\alpha}{2}} \right] e^{(1+\alpha) \int_{T_0}^t Q_2 ds}.$$

We treat the case $T_1 \leq t \leq \min\{T, T_e\}$. Using Lemma 4.5 with $\gamma = \frac{\alpha}{2}$, and Lemma 6.9 we get

$$\begin{aligned} e^{2A} |b| &\leq e^{2A} X_1^{1-\alpha} [M_{\widehat{D}}(t)(1 + |\log d(\mathbf{X})|) + X_1^{\frac{\alpha}{2}} d(\mathbf{X})^{-1+\frac{\alpha}{2}}] \\ &\leq g e^{(1+\alpha) \int_{T_0}^s Q_1 ds} X_1(T_0)^{1-\alpha} [M(1 + |\log d(\mathbf{X})|) + \delta_1^{\frac{\alpha}{2}} d(\mathbf{X})^{-1+\frac{\alpha}{2}}]. \end{aligned}$$

Also recall that $M_{\widehat{D}}(t) \leq M$. To integrate this bound from T_0 to t we split into two integrals from T_0 to T_1 and T_1 to t . For $t \in [T_0, T_1]$ we can estimate the factor in square brackets independent of t using $d(\mathbf{X}) \geq \frac{1}{2}\delta_2$:

$$g X_1(T_0)^{1-\alpha} [M(1 + |\log \delta_2|) + \delta_1^{\frac{\alpha}{2}} \delta_2^{-1+\frac{\alpha}{2}}] \int_{T_0}^{T_1} e^{(1+\alpha) \int_{T_0}^s Q_1 d\tau} ds.$$

The remaining integral can be estimated as follows:

$$\begin{aligned} \int_{T_0}^{T_1} e^{(1+\alpha) \int_{T_0}^s Q_1 d\tau} ds &\leq \int_{T_0}^t \frac{Q_1}{Q_1} e^{(1+\alpha) \int_{T_0}^s Q_1 d\tau} ds \leq \beta^{-1} (1+\alpha)^{-1} e^{(1+\alpha) \int_{T_0}^s Q_1 d\tau} \Big|_{s=T_0}^{s=t} \\ &\lesssim e^{(1+\alpha) \int_{T_0}^t Q_1 d\tau}. \end{aligned}$$

Hence for sufficiently small δ_2

$$\int_{T_0}^{T_1} e^{2A} |b| \, ds \lesssim g X_1(T_0)^{1-\alpha} [M |\log \delta_2| + \delta_1^{\frac{\alpha}{2}} \delta_2^{-1+\frac{\alpha}{2}}] e^{(1+\alpha) \int_{T_0}^{T_1} Q_1}.$$

For the remaining part $\int_{T_1}^t e^{2A} |b| \, ds$, we use Proposition 6.8 again, and find the bound for small δ_2

$$\begin{aligned} & g X_1(T_0)^{1-\alpha} \int_{T_1}^t e^{(1+\alpha) \int_{T_0}^s Q_1 ds} [M(1 + |\log d(\mathbf{X})|) + \delta_1^{\frac{\alpha}{2}} d(\mathbf{X})^{-1+\frac{\alpha}{2}}] \, ds \\ & \leq g e^{(1+\alpha) \int_{T_0}^t Q_1 ds} X_1(T_0)^{1-\alpha} \int_{T_1}^t [M(1 + |\log d(\mathbf{X})|) + \delta_1^{\frac{\alpha}{2}} d(\mathbf{X})^{-1+\frac{\alpha}{2}}] \, ds \\ & \lesssim g X_1(T_0)^{1-\alpha} [M |\log \delta_2| + \delta_1^{\alpha/2} \delta_2^{-1+\alpha/2}] e^{(1+\alpha) \int_{T_0}^t Q_1 ds}. \end{aligned}$$

Using the second estimate from Lemma 6.9 the claim follows for the case $T_1 \leq t \leq \min\{T, T_e\}$. The calculation for $t \leq T_1$ is similar (and slightly simpler).

Next, using again Lemma 4.5 with $\gamma_1 = \gamma_2 = \alpha/2$ and Lemma 6.9,

$$\begin{aligned} e^{-A} |c| & \leq e^{-A} X_2^{1-\alpha} [M + X_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2}] \\ & \leq g e^{-\int_{T_0}^t Q_2 \, ds} X_2(T_0)^{1-\alpha} [M + \delta_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2}]. \end{aligned}$$

Hence

$$\begin{aligned} \int_{T_0}^t e^{-A} |c| \int_{T_0}^s e^{2A} |b| \, ds & \lesssim g X_2(T_0)^{1-\alpha} X_1(T_0)^{1-\alpha} [M |\log \delta_2| + \delta_1^{\alpha/2} \delta_2^{-1+\alpha/2}] \\ & \quad \times \int_{T_0}^t e^{\int_{T_0}^s Q_2 \, d\tau} [M + \delta_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2}] \, ds. \end{aligned}$$

We continue to estimate the last integral:

$$\begin{aligned} & \int_{T_0}^t e^{\int_{T_0}^s Q_2 \, d\tau} [M + \delta_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2}] \, ds \\ & = M \int_{T_0}^t e^{\int_{T_0}^s Q_2 \, d\tau} \, ds + \int_{T_0}^t e^{\int_{T_0}^s Q_2 \, d\tau} \delta_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2} \, ds \\ & \leq M \int_{T_0}^t \frac{Q_2}{Q_2} e^{\int_{T_0}^s Q_2 \, d\tau} \, ds + e^{\int_{T_0}^t Q_2 \, d\tau} \delta_1^{\alpha/2} \int_{T_0}^t d(\mathbf{X})^{-1+\alpha/2} \, ds \\ & \lesssim e^{\int_{T_0}^t Q_2 \, ds} [M + \delta_1^{\alpha/2} \delta_2^{-1+\alpha/2}] \end{aligned}$$

where we used the familiar splitting at T_1 . Thus, finally we get

$$\begin{aligned} \int_{T_0}^t e^{-A} |c| \int_{T_0}^s e^{2A} |b| \, ds & \lesssim g X_1(T_0)^{1-\alpha} [M |\log \delta_2| + \delta_1^{\alpha/2} \delta_2^{-1+\alpha/2}] \\ & \quad \times [M + \delta_1^{\alpha/2} \delta_2^{-1+\alpha/2}] e^{\int_{T_0}^t Q_2 \, ds} X_2(T_0)^{1-\alpha}. \end{aligned}$$

It remains to apply key Lemma 6.7 to estimate the factor $e^{\int_{T_0}^t Q_2 \, ds} X_2(T_0)^{1-\alpha}$, which is less than

$$\delta_2^{1-\alpha} e^{\int_{T_0}^t Q_2 \, ds} e^{-(1-\alpha) \int_{T_0}^t Q_2 \, ds} \leq \delta_2^{1-\alpha} e^{\alpha \int_{T_0}^t Q_2 \, ds} = \delta_2^{1-2\alpha} \delta_2^{\alpha} e^{\alpha \int_{T_0}^t Q_2 \, ds}.$$

Now observe that the expression $\left[M|\log \delta_2| + \delta_1^{\alpha/2} \delta_2^{-1+\alpha/2} \right] [M + \delta_1^{\alpha/2} \delta_2^{-1+\alpha/2}] \delta_2^{1-2\alpha}$ is a harmless factor since $\alpha < 1/4$.

□

Lemma 6.12. *For sufficiently small δ_2 and $t \in [T_0, \min\{T, T_e\}]$ we have the following inequalities.*

$$(41) \quad v_1(t) \leq g \left[Q_2 + M X_2^{1-\alpha} |\log d(\mathbf{X})| + \delta_2^{1-\alpha} d(\mathbf{X})^{-1+\alpha} \right] e^{-\int_{T_0}^t Q_2 ds},$$

$$(42) \quad v_2(t) \leq g X_1(T_0)^{1-\alpha} \left[M|\log d(\mathbf{X})| + \delta_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2} \right] e^{\alpha \int_{T_0}^t Q_2 ds},$$

$$(43) \quad \int_{T_0}^t v_1 e^{\alpha \int_{T_0}^s Q_2 d\tau} ds \leq g$$

$$(44) \quad \int_{T_0}^t v_2 ds \leq g e^{\alpha \int_{T_0}^t Q_2 ds}$$

$$(45) \quad \int_{T_0}^t v_2 e^{\alpha \int_{T_0}^s Q_2 d\tau} ds \leq g e^{2\alpha \int_{T_0}^t Q_2 ds},$$

$$(46) \quad \int_{T_0}^t v_1 f_1 ds \leq g \delta_1^{1-\alpha} \delta_2^\alpha,$$

$$(47) \quad \int_{T_0}^t v_2 f_1 ds \leq g \delta_1^{1-\alpha} \delta_2^{\alpha/2} X_1(T_0)^{1-\alpha} e^{2\alpha \int_{T_0}^t Q_2 d\tau},$$

$$(48) \quad \int_{T_0}^t v_1 f_2 ds \leq g X_2(T_0)^{1-\alpha},$$

where $g = g(\alpha, \beta, \delta, M)$ is a harmless factor.

Proof. The estimates (41)-(44) follow from Lemma 4.5, Lemma 6.9, Proposition 6.8 and the usual splitting of the interval of integration into $[T_0, T_1]$ and $[T_1, \min\{T, T_e\}]$. (45) follows easily from (44) and Lemma 6.9.

Using Lemma 6.11 and Lemma 4.5 we get

$$v_1 f_1(s) \leq g \delta_1^{1-\alpha} \delta_2^\alpha \left[Q_2 + M X_2^{1-\alpha} |\log d(\mathbf{X})| + \delta_2^{1-\alpha} d(\mathbf{X})^{-1+\alpha} \right] e^{(-1+\alpha) \int_{T_0}^s Q_2 d\tau}.$$

Note how the exponential growth of the factor f_1 was cancelled by the exponential factor in v_1 . By integrating, we get:

$$\begin{aligned} \int_{T_0}^t v_1 f_1 ds &\leq g \delta_1^{1-\alpha} \delta_2^\alpha \int_{T_0}^t e^{(-1+\alpha) \int_{T_0}^s Q_2 d\tau} \left[Q_2 + M \delta_2^{1-\alpha} |\log d(\mathbf{X})| + \delta_2^{1-\alpha} d(\mathbf{X})^{-1+\alpha} \right] ds \\ &\leq g \delta_1^{1-\alpha} \delta_2^\alpha \left[M \delta_2^{1-\alpha} |\log \delta_2| + 1 \right] \end{aligned}$$

giving (46) since the factor in square brackets is harmless and can be absorbed into g .

Proceeding analogously to prove (47) we find

$$v_2 f_1(s) \leq g \delta_1^{1-\alpha} \delta_2^\alpha X_1(T_0)^{1-\alpha} e^{2\alpha \int_{T_0}^s Q_2 d\tau} \left[M|\log d(\mathbf{X})| + \delta_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2} \right]$$

which after integration from T_0 to t can be estimated as follows:

$$\begin{aligned} \int_{T_0}^t v_2 f_1 \, ds &\leq g \delta_1^{1-\alpha} \delta_2^\alpha X_1(T_0)^{1-\alpha} e^{2\alpha \int_{T_0}^s Q_2 \, d\tau} \left[M |\log \delta_2| + \delta_1^{\alpha/2} \delta_2^{-1+\alpha/2} \right] \\ &\leq g \delta_1^{1-\alpha} \delta_2^{\alpha/2} X_1(T_0)^{1-\alpha} e^{2\alpha \int_{T_0}^t Q_2 \, d\tau} \left[M \delta_2^{\alpha/2} |\log \delta_2| + \delta_1^{\alpha/2} \delta_2 \right]. \end{aligned}$$

Again the factor in square brackets can be absorbed into g giving (47).

(48) is derived using the same techniques. □

Lemma 6.13. *Along a particle trajectory, for $T_0 \leq t \leq \min\{T, T_e\}$,*

$$(49) \quad (Hf_0)(t) \leq g \|f_0\|_\infty(t) e^{\alpha \int_{T_0}^t Q_2 \, ds}$$

holds, where $\|f_0\|_\infty(t) = \sup\{|f_0(s)| : s \in [T_0, t]\}$.

Proof. Using Lemmas 6.11, 6.12, we get

$$(50) \quad f_0 + f_1 \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 f_0 \leq g \|f_0\|_\infty(t) e^{\alpha \int_{T_0}^t Q_2 \, ds}.$$

Recall that products and exponentials of harmless factors are harmless, too.

Next, using Lemma 6.10, 6.11 and 6.12

$$\begin{aligned} f_2 + f_1 \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 f_2 &\leq g X_2(T_0)^{1-\alpha} e^{\alpha \int_{T_0}^t Q_2 \, ds} \\ &= g X_2(T_0)^{1-2\alpha} X_2(T_0)^\alpha e^{\alpha \int_{T_0}^t Q_2 \, ds} \\ &\leq g X_2(T_0)^{1-2\alpha} \delta_2^\alpha \end{aligned}$$

with the key Lemma 6.7 in the last step to cancel of $e^{\alpha \int_{T_0}^t Q_2 \, ds}$ using the factor $X_2(T_0)^\alpha$.

So for $v_2 \left[f_2 + f_1 \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 f_2 \right]$ we obtain the upper bound

$$\begin{aligned} v_2 g X_2(T_0)^{1-2\alpha} \delta_2^\alpha &\leq g e^{\alpha \int_{T_0}^t Q_2} X_2(T_0)^{1-2\alpha} \delta_2^\alpha X_1(T_0)^{1-\alpha} \left[M |\log d(\mathbf{X})| + \delta_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2} \right] \\ &\leq g \delta_2^{2\alpha} X_2(T_0)^{1-3\alpha} \delta_1^{1-\alpha} \left[M |\log d(\mathbf{X})| + \delta_1^{\alpha/2} d(\mathbf{X})^{-1+\alpha/2} \right] \end{aligned}$$

using the key Lemma 6.7 again to cancel $e^{\alpha \int_{T_0}^t Q_2}$. Thus we see that

$$\exp \left(\int_{T_0}^t v_2 \left[f_2 + f_1 \exp \left(\int_{T_0}^s v_1 f_1 \right) \int_{T_0}^s v_1 f_2 \right] \right) \leq g$$

Finally, by (50) and Lemma 6.12

$$\begin{aligned} \int_{T_0}^t v_2 \left[f_0 + f_1 \exp \left(\int_{T_0}^s v_1 f_1 \right) \int_{T_0}^s v_1 f_0 \right] &\leq \\ g \|f_0\|_\infty(t) \int_{T_0}^t v_2 e^{\alpha \int_{T_0}^s Q_2 \, d\tau} \, ds &\leq g \|f_0\|_\infty(t) e^{2\alpha \int_{T_0}^t Q_2 \, ds}. \end{aligned}$$

Thus, in total we get

$$(51) \quad \begin{aligned} f_0 + f_1 \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 f_0 &\leq g \|f_0\|_\infty(t) e^{\alpha \int_{T_0}^t Q_2 ds} \\ f_2 + f_1 \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 f_2 &\leq g X_2(T_0)^{1-2\alpha} \delta_2^\alpha \end{aligned}$$

$$(52) \quad \exp \left(\int_{T_0}^t v_2 \left[f_2 + f_1 \exp \left(\int_{T_0}^s v_1 f_1 \right) \int_{T_0}^s v_1 f_2 \right] \right) \leq g$$

$$(53) \quad \int_{T_0}^t v_2 \left[f_0 + f_1 \exp \left(\int_{T_0}^s v_1 f_1 \right) \int_{T_0}^s v_1 f_0 \right] \leq g \|f_0\|_\infty(t) e^{2\alpha \int_{T_0}^t Q_2 ds}.$$

Note that

$$(51) \times (53) \leq g X_2(T_0)^{1-4\alpha} \delta_2^{4\alpha} \|f_0\|_\infty(t) \leq g$$

using again the key Lemma to get rid of the factor $e^{2\alpha \int_{T_0}^t Q_2}$, and in the very last step we used $\alpha \in (0, 1/4)$ and Lemma 6.10. Combining the inequalities (51)-(53) gives

$$\begin{aligned} &\left(f_2 + f_1 \exp \left(\int_{T_0}^t v_1 f_1 \right) \int_{T_0}^t v_1 f_2 \right) \\ &\times \exp \left(\int_{T_0}^t v_2 \left[f_2 + f_1 \exp \left(\int_{T_0}^s v_1 f_1 \right) \int_{T_0}^s v_1 f_2 \right] \right) \\ &\times \int_{T_0}^t v_2 \left[f_0 + f_1 \exp \left(\int_{T_0}^s v_1 f_1 \right) \int_{T_0}^s v_1 f_0 \right] \\ &\leq g X_2(T_0)^{1-4\alpha} \delta_2^{4\alpha} \|f_0\|_\infty(t) \leq g \|f_0\|_\infty(t) \end{aligned}$$

In view of (28), (49) now follows. $\square$

6.4. Proof of the main technical theorem.

Proof of theorem 6.3. At time $t = T$, any $x \in D$ is occupied by a particle, i.e. $x = \mathbf{X}(T)$ for some particle trajectory. Let us write

$$\partial_{x_j} \omega(\mathbf{X}(t), t) = \xi_j(t)$$

along that particle trajectory, and so by (31),

$$|\xi_1(t)| \leq (Hf_0)(t) + e^A \int_{T_0}^t |a| e^{-A} (Hf_0)(s) ds = (Hf_0)(t) + e^A \int_{T_0}^t v_1 (Hf_0)(s) ds.$$

First note that by Lemmas 6.13, 6.10, 6.12 and 6.9

$$\begin{aligned} e^A \int_{T_0}^t v_1(s) (Hf_0)(s) ds &\leq e^A g \|f_0\|_\infty(t) \int_{T_0}^t v_1 e^{\alpha \int_{T_0}^s Q_2 d\tau} ds \leq g R X_2(T_0)^{1-\alpha} e^A \\ &\leq g R e^{-(1-\alpha) \int_{T_0}^t Q_2 ds} e^{\int_{T_0}^t Q_2 ds} \leq g R e^{\alpha \int_{T_0}^t Q_2 ds}. \end{aligned}$$

Moreover again by Lemmas 6.13, 6.10

$$(Hf_0)(t) \leq g R e^{\alpha \int_{T_0}^t Q_2 ds}.$$

This results in

$$(54) \quad |\xi_1(t)| \leq g R e^{\alpha \int_{T_0}^t Q_2 ds}.$$

Next we estimate $|\xi_1(t)|X_1(t)^\alpha$. First we use (54) and insert $X_1(t)$ from (34). Then Lemma 6.9 allows us to replace Q_1 by Q_2 in the one of the arguments of the exponential function, so we get the estimate

$$(55) \quad \begin{aligned} |\xi_1(t)|X_1(t)^\alpha &\leq g R X_1(t)^\alpha e^{\alpha \int_{T_0}^t Q_1 ds} \leq g R \delta_1^\alpha e^{-\alpha \int_{T_0}^t Q_1 ds} e^{\alpha \int_{T_0}^t Q_2 ds} \\ &\leq g R \delta_1^\alpha e^{-\alpha \int_{T_0}^t Q_2 ds} e^{\alpha \int_{T_0}^t Q_2 ds} \leq g R \delta_2^\alpha \end{aligned}$$

since $\delta_1 < \delta_2$. In fact, this was the most critical estimate in the whole proof of the main result, since the dangerous factor e^A was barely cancelled.

We now derive a similar estimate for $|\xi_2(t)|X_2(t)^\alpha$. From the second line of (31) and the assumptions on initial conditions,

$$(56) \quad |\xi_2(t)| \leq R e^{-A} + e^{-A} \int_{T_0}^t v_2 H f_0 ds + e^{-A} \int_{T_0}^t e^{2A} |b| \int_{T_0}^s v_1 H f_0 ds.$$

By Lemma 6.13 and 6.10 $H f_0(s)$ has the upper bound

$$R g X_2(T_0)^{1-\alpha} e^{\alpha \int_{T_0}^s Q_2 d\tau}.$$

Therefore the second summand can be estimated as follows

$$\begin{aligned} e^{-A} \int_{T_0}^t v_2 H f_0 ds &\leq e^{-A} R g \int_{T_0}^t v_2 e^{\alpha \int_{T_0}^s Q_2 d\tau} ds \leq R g e^{-A} e^{2\alpha \int_{T_0}^t Q_2 ds} \\ &\leq R g e^{(-1+2\alpha) \int_{T_0}^t Q_2 ds} \leq g R, \end{aligned}$$

where we also used $X_2(T_0) \leq \delta_2$, (45), Lemma 6.9 and $\alpha < 1/4$.

For the third summand of (56) we use the upper bound for $H f_0$ again:

$$\begin{aligned} e^{-A} \int_{T_0}^t e^{2A} |b| \int_{T_0}^s v_1 H f_0 ds &\leq g R X_2(T_0)^{1-\alpha} e^{-A} \int_{T_0}^t e^{2A} |b| \int_{T_0}^s v_1 e^{\alpha \int_{T_0}^\tau Q_2 d\zeta} d\tau ds \\ &\leq g R \delta_2^{1-\alpha} e^{-A} \int_{T_0}^t e^{2A} |b| ds. \end{aligned}$$

In the last estimate (43) was used. Now note that by (40)

$$\int_{T_0}^t e^{2A} |b| ds \leq g e^{(1+\alpha) \int_{T_0}^t Q_2 ds}.$$

After combining this with the e^{-A} factor we see that we can bound the third summand by $g R$, i.e. $|\xi_2(t)| \leq g R$. This means that

$$(57) \quad |\xi_2(t)|X_2^\alpha \leq g R \delta_2^\alpha.$$

The inequalities (57) and (55) imply

$$M_D(T) \leq g(\alpha, \beta, \boldsymbol{\delta}, M) R \delta_2^\alpha + \|\omega\|_\infty =: \mathcal{N}(R, \alpha, \beta, \boldsymbol{\delta}, M).$$

It remains to show that $\mathcal{N}$ is a harmless nonlinearity. Therefore, let α, β, R be given. Recall that g has the property that $g(\alpha, \beta, \delta_2^p, \delta_2, R)$ is bounded as $\delta_2 \rightarrow 0$ for all $p > p_0$ with some $p_0 > 0$. Hence

$$g(\alpha, \beta, \delta_2^p, \delta_2, R)R\delta_2^\alpha + \|\omega\|_\infty < R$$

for sufficiently small $\delta_2 > 0$ and $R > \|\omega\|_\infty$. $\square$

7. PROOF OF THE MAIN RESULT

Proof of Theorem 1.4. Let $\alpha \in (0, \frac{1}{4})$, $0 < \delta_3 < 1/2$ and $R > \|\omega\|_\infty$ be a given nonnegative number. Fix small positive δ_1, δ_2 such that the following set of inequalities hold true:

$$(58) \quad \delta_1, \delta_2 \leq \rho, \quad \beta_0 - A|\delta|^{1-\alpha}R \geq \frac{1}{2}\beta_0.$$

$$(59) \quad M_D(0) < R, \quad \left| \frac{\partial \omega_0(x)}{\partial x_1} \right| \leq R x_2^{1-\alpha}, \quad \left| \frac{\partial \omega_0(x)}{\partial x_2} \right| \leq R \quad (x \in D)$$

where A, β_0, ρ are the numbers from the Definition 1.2 of the hyperbolicity of the flow. Note that the box can be chosen so small that that (59) holds. This a consequence of $\frac{\partial \omega_0}{\partial x_2}(0, x_2) = \frac{\partial \omega_0}{\partial x_1}(x_1, 0) = 0$ and the C^2 -smoothness of ω_0 .

Claim: If the box $\widehat{D}$ satisfies the controlled feeding conditions with parameter R , then we have the bound

$$(60) \quad M_D(t) \leq R \quad (t \in [0, \infty)).$$

Assume (60) is not true for all times, i.e. there is a time $\widetilde{T}$ such that $M_D(\widetilde{T}) > R$. Since the solution ω is sufficiently smooth in time by assumption, $M_D(t)$ is a continuous function of t . Because $M_D(0) < R$, by the intermediate value theorem, there exists a time $T \in (0, \widetilde{T})$ such that $M_D(t) < R$ holds on $[0, T)$ and $M_D(T) = R$. Observe also that automatically $M_{\widehat{D}}(t) \leq R$ for $t \leq T$.

Now note that by (58), the flow is $\frac{1}{2}\beta_0$ -hyperbolic in the box D on the time interval $[0, T]$. This can be seen as follows. Because of (17) and the feeding conditions, $M(x, t) \leq M_{\widehat{D}}(t) \leq R$ for all $x \in D$ and $t \in [0, T]$. Thus by (58)

$$Q_i(x, t) = Q_i(x, t) + A|x|^{1-\alpha}M(x, t) - A|x|^{1-\alpha}M(x, t) \geq \beta_0 - \frac{1}{2}\beta_0.$$

Upon shrinking δ_1 and δ_2 further (if necessary) and using (59) we can achieve that the assumptions of Theorem 6.3 are satisfied, the arguments in Section 6 hold and for the harmless nonlinear function $\mathcal{N}$ from theorem 6.3 the following inequality is true:

$$(61) \quad \mathcal{N}(R, \alpha, \frac{1}{2}\beta_0, \delta_1, \delta_2, R) < R.$$

From now on δ is fixed.

On the one hand, on $[0, T]$, we have $M_D(t) \leq R$. But applying Theorem 6.3 with $K = R$ and (61), we get

$$M_D(T) \leq \mathcal{N}(R, \alpha, \frac{1}{2}\beta_0, \delta_1, \delta_2, R) < R,$$

a contradiction. This proves our claim (60).

Now we prove the exponential bound on the gradient growth. At an arbitrary time $t \geq 0$, each $x \in D$ is occupied by a particle $\mathbf{X}(t)$ that has entered the box at some earlier

time T_0 , or $T_0 = 0$ if the particle started in D at $t = 0$. The same calculation leading to (55) gives

$$\left| \frac{\partial \omega}{\partial x_1}(\mathbf{X}(t), t) \right| \leq g R e^{\alpha \int_{T_0}^t Q_2 ds}.$$

for all $t \in [T_0, T_e]$. We apply now Lemma 4.4:

$$\int_{T_0}^t Q_2 ds \lesssim (\|\omega\|_\infty + R(|\boldsymbol{\delta}| + \delta_3)^{1-\alpha})(t - T_0) + \|\omega\|_\infty \int_{T_0}^t |\log d(\mathbf{X})| ds.$$

The integral containing the logarithmic term can be estimated using the familiar splitting at T_1 and gives

$$\int_{T_0}^t |\log d(\mathbf{X})| ds \lesssim |\log \delta_2|(t - T_0) + |\log \delta_2|.$$

Thus, finally,

$$\left| \frac{\partial \omega}{\partial x_1}(\mathbf{X}(t), t) \right| \leq C(\alpha, \beta, \boldsymbol{\delta}, \delta_3, R, \|\omega\|_\infty) e^{\alpha(C\|\omega\|_\infty + R(|\boldsymbol{\delta}| + \delta_3)^{1-\alpha} + |\log \delta_2|)t}.$$

The derivative in x_2 -direction is bounded by gR as seen in the proof of Theorem 6.3. This concludes the proof of Theorem 1.4. $\square$

Remark 7.1. Our main result remains valid if instead of $\omega_0 \in C^2$ we only assume $\omega_0 \in C^{1,\gamma}$ with $\gamma \in (3/4, 1)$ (note that (59) is still true in this case). Recall that in [11] a solution in $C^{1,\gamma}$ was constructed such that the gradient growth close to the origin is at least exponential. This allows us to state the following interesting *conditional result*:

Corollary 7.2. Suppose a solution ω in $C^{1,\gamma}$ as in [11] exists that satisfies the feeding conditions in some box. Then exponential gradient growth near the origin is optimal in the class of $C^{1,\gamma}$ -solutions.

Remark 7.3. At this point, we address the difficulties in applying the techniques developed here to the case of time-dependent feeding. Assume for the sake of the discussion that in (4) we replace the constant R by

$$R(t) = R_1 t + R_2$$

with positive constants R_1, R_2 . As a direct consequence, f_0 grows in time. The corresponding inequality in Lemma 6.10 will roughly read

$$f_0(t) \lesssim gR(T_0)X_2(T_0)^{1-\alpha}.$$

This produces, for example, via (49) a growth in our bounds for $M_{\hat{D}}(t)$, so that $M_{\hat{D}}(t)$ cannot be bounded by a time-independent constant anymore. The consequences are as follows:

- The hyperbolicity condition (6) is no longer sufficient to stabilize the flow. It has to be considerably strengthened, for example by requiring the flow to be at least β -hyperbolic from the outset. Lemma 6.9 does not seem to go through, so one may have to strengthen the condition even further, e.g. by requiring

$$(62) \quad a(x, t) \geq \beta > 0.$$

This, however, has the disadvantage that we have no sufficient condition on the initial data to justify the validity of (62).

- A destabilizing effect is also felt in all estimates of Lemma 6.12, especially when the now time-dependent bound for $M_{\widehat{D}}(t)$ is integrated along a particle trajectory with a growing factor (e.g. $\exp(\alpha \int_{T_0}^t Q_s d\tau)$). This leads to worse estimates for the quantities f_1, f_2 and hence for Hf_0 , which are a vital part of the main argument.

In this paper, we leave the problem of finding a suitable treatment for time-dependent feeding open.

8. ACKNOWLEDGEMENTS

The authors cordially thank A. Kiselev for suggesting the problem and a great number of helpful discussions, as well the anonymous reviewers for their careful reading of the manuscript and helpful comments. VH would like to express his gratitude to the Deutsche Forschungsgemeinschaft (German Research Foundation), without whose financial support (FOR HO 5156/1-1 and FOR HO 5156/1-2) the present research could not have been undertaken. VH also acknowledges partial support by NSF grant nsf-dms 1412023.

9. APPENDIX

9.1. **Appendix A.** We use the Kronecker symbol

$$\delta_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}.$$

Proposition 9.1. *For all $x, y \in [0, 1]^2$, $x \neq y$ the following estimates hold.*

$$\begin{aligned} |G_i^k(x, y)| &\lesssim |y - x|^{-1} x_i^{-1} \\ \left| \frac{\partial G_i^k}{\partial x_j}(x, y) \right| &\lesssim |y - x|^{-3} \end{aligned}$$

($i, k = 1, 2$).

The proofs are straightforward calculations based on the identities in Appendix B, and the reflection inequalities:

$$(63) \quad |y - \widetilde{x}| \geq |y - x|, \quad |y - \bar{x}| \geq |y - x|, \quad |y + x| \geq |y - \bar{x}|, \quad |y + x| \geq |y - \widetilde{x}|$$

holding for $x, y \in [0, 1]^2$. Also, use the obvious inequalities

$$x_2 \leq |y - \bar{x}|, x_1 \leq |y - \widetilde{x}|.$$

We observe some useful relations for the kernels G_i^k and their derivatives. Let G stand for any G_i^k and let

$$\Omega_x = (-x_1, 1 - x_1) \times (-x_2, 1 - x_2).$$

G has the form $G(x, y) = \widetilde{G}(y - x, x, y)$, where $\widetilde{G}(z, \eta, \mu)$ is smooth provided $\eta, \mu \in (0, 1)^2, z \in \Omega_x \setminus \{0\}$. For example, if $G = G_1^1$ then

$$\widetilde{G}(z, \eta, \mu) = \frac{\mu_1 z_1}{|z|^2 |\mu - \widetilde{\eta}|^2}.$$

Note that for $x \neq y, x, y \in (0, 1)^2$,

$$(64) \quad \begin{aligned} (\partial_{x_j} G)(x, y) &= (\partial_{\eta_j} \tilde{G})(y - x, x, y) - (\partial_{z_j} \tilde{G})(y - x, x, y) \\ (\partial_{y_j} G)(x, y) &= (\partial_{\mu_j} \tilde{G})(y - x, x, y) + (\partial_{z_j} \tilde{G})(y - x, x, y) \end{aligned}$$

so that

$$(\partial_{x_j} G)(x, y) = -(\partial_{y_j} G)(x, y) + (\partial_{\eta_j} \tilde{G})(y - x, x, y) + (\partial_{\mu_j} \tilde{G})(y - x, x, y).$$

Moreover, we always have

$$(65) \quad |\tilde{G}(z, x, y)|, \left| \frac{\partial \tilde{G}}{\partial \eta_j}(z, x, x + z) \right|, \left| \frac{\partial \tilde{G}}{\partial \mu_j}(z, x, x + z) \right| \leq C(\eta) |z|^{-1}.$$

where $C(\eta)$ is uniformly bounded if η varies in a compact subset of $(0, 1)^2$.

Proposition 9.2.

$$\frac{\partial G_i^k}{\partial x_j} = -\frac{\partial G_i^k}{\partial y_j} + x_i^{-2} \delta_{ij} \mathcal{O}(|y - x|^{-1})$$

Proof. This is a tedious, but straightforward estimation using the identities of Appendix B and the reflection inequalities. $\square$

Proposition 9.3 (Derivatives of Q_i).

$$\begin{aligned} \frac{\partial Q_i}{\partial x_j} &= c_0 P.V. \int_{[0,1]^2} \left[\frac{\partial G_i^1}{\partial x_j} + \frac{\partial G_i^2}{\partial x_j} \right] \omega(y) dy \\ &\quad - \omega(x) \lim_{\delta \rightarrow 0^+} \int_{\partial B(x, \delta)} G_i^r \cdot \nu_j d\sigma + \frac{\partial Q_i^r}{\partial x_j} \end{aligned}$$

Proof. Write $(G_i^1 + G_i^2)(x, y) := G(x, y)$. G has again the form $G(x, y) = \tilde{G}(y - x, x, y)$, where $\tilde{G}(z, \eta, \mu)$ is smooth provided $\eta, \mu \in (0, 1)^2, z \in \Omega_x \setminus \{0\}$. Also (64), (65) hold for $\tilde{G}$. Now

$$(66) \quad \begin{aligned} \frac{\partial}{\partial x_j} \int_{\Omega_x} \tilde{G}(z, x, x + z) \omega(x + z) dz &= \int_{\Omega_x} \tilde{G}(z, x, x + z) \frac{\partial \omega}{\partial z_j}(x + z) dz \\ &\quad + \int_{\Omega_x} \partial_{x_j} (\tilde{G}(z, x, x + z)) \omega(x + z) dz \\ &\quad - \int_{\partial \Omega_x} \tilde{G}(z, x, x + z) \omega(x + z) \nu_j d\sigma \end{aligned}$$

where ν_j denotes the j -th component of the unit outer normal. This is a standard differentiation result (note the bounds (65)).

Now consider the integral in the line (66), exclude the singularity and integrate by parts:

$$(67) \quad \int_{\Omega_x} \tilde{G}(z, x, x+z) \frac{\partial \omega}{\partial z_j}(x+z) dz = - \int_{\Omega_x \setminus B(0, \delta)} \partial_{z_j}(\tilde{G}(z, x, x+z)) \omega(x+z) dz \\ + \int_{\partial \Omega_x} \tilde{G}(z, x, x+z) \omega(x+z) \nu_j d\sigma \\ - \int_{\partial B(0, \delta)} \tilde{G}(z, x, x+z) \omega(x+z) \nu_j d\sigma$$

Observe that by (64),

$$-\partial_{z_j}(\tilde{G}(z, x, x+z)) + \partial_{x_j}(G(z, x, x+z)) = (\partial_{x_j} G)(z, x, x+z).$$

So combining (66) and (67), we finally get

$$\frac{\partial}{\partial x_j} \int_{\Omega_x} \tilde{G}(z, x, x+z) \omega(x+z) dz = - \int_{\partial B(0, \delta)} \tilde{G}(z, x, x+z) \omega(x+z) \nu_j d\sigma \\ + \int_{\Omega_x} (\partial_{x_j} G)(z, x, x+z) \omega(x+z) dz + \int_{B(0, \delta)} \partial_{x_j}(\tilde{G}(z, x, x+z)) \omega(x+z) dz.$$

Replacing $x+z$ by y and sending $\delta \rightarrow 0$ yields the statement. $\square$

We define

$$d_1(x) := \min\{x_1, x_2\}$$

which is the distance of the point x to the coordinate axes. Observe also that

$$(68) \quad \frac{1}{2}x_r \leq y_r \leq \frac{3}{2}x_r$$

for $y \in B(x, \frac{1}{2}d_1(x))$, $r = 1, 2$. For the entire appendix, we shall write that $M = M_{\widehat{D}}$, i.e.

$$\left| \frac{\partial \omega}{\partial x_j}(x) \right| \leq M x_j^{-\alpha} \quad (x \in \widehat{D}, j = 1, 2)$$

holds, implying also the inequalities

$$(69) \quad |\omega(x)| \lesssim M x_j^{1-\alpha} \quad (x \in \widehat{D}, j = 1, 2)$$

(by the fact that ω vanishes identically on the coordinate axes).

Proposition 9.4. *For $i \neq j$, we have*

$$\left| \frac{\partial(G_i^1 + G_i^2)}{\partial x_j} \right| \lesssim x_i^{-\gamma_1 - \gamma_2} x_j^{\gamma_2} |y - x|^{-(3-\gamma_1)}.$$

where $\gamma_1, \gamma_2 \in \mathbb{R}$, $0 \leq \gamma_1 + \gamma_2 \leq 1$.

Proof. We do the proof in the case $i = 2, j = 1$, the other case being analogous. The proof of the proposition is based on a cancellation property of the kernels G_2^1 and G_2^2 and

requires quite tedious computations. First calculate the sum of $\partial_{x_1} G_2^2$ and $\partial_{x_1} G_2^1$ and see that it can be grouped into three the expressions

$$\begin{aligned} \frac{2y_2(y_1 - x_1)^2}{|y - x|^4|y - \bar{x}|^2} - \frac{2y_2(y_1 + x_1)^2}{|y + x|^2|y - \tilde{x}|^4} &= (A) \\ \frac{2y_2(y_1 - x_1)^2}{|y - x|^2|y - \bar{x}|^4} - \frac{2y_2(y_1 + x_1)^2}{|y - \tilde{x}|^2|y + x|^4} &= (B) \\ \frac{y_2}{|y - \tilde{x}|^2|y + x|^2} - \frac{y_2}{|y - x|^2|y - \bar{x}|^2} &= (C) \end{aligned}$$

These can be further written as

$$\begin{aligned} &\frac{2y_2(y_1 - x_1)^2}{|y - \bar{x}|^2} [|y - x|^{-4} - |y - \tilde{x}|^{-4}] + \frac{2y_2}{|y - \tilde{x}|^4} \left(\frac{(y_1 - x_1)^2}{|y - \bar{x}|^2} - \frac{(y_1 + x_1)^2}{|y + x|^2} \right) \\ &= (1) + (2) \\ &\frac{2y_2}{|y - \bar{x}|^4} \left[\frac{(y_1 - x_1)^2}{|y - x|^2} - \frac{(y_1 + x_1)^2}{|y - \tilde{x}|^2} \right] + \frac{2y_2}{|y - \tilde{x}|^2} \left[\frac{(y_1 + x_1)^2}{|y - \bar{x}|^4} - \frac{(y_1 + x_1)^2}{|y + x|^4} \right] \\ &= (3) + (4) \\ &\frac{y_2}{|y + x|^2} [|y - \tilde{x}|^{-2} - |y - x|^{-2}] + \frac{y_2}{|y - x|^2} [|y + x|^{-2} - |y - \bar{x}|^{-2}] \\ &= (5) + (6) \end{aligned}$$

Let us estimate expression (1). Using $|y - \tilde{x}|^2 - |y - x|^2 = 4x_1y_1$ and the relations $y_2 \leq |y - \bar{x}|$, $(y_1 - x_1)^2 \leq |y - x|^2$, $y_1 \leq (y_1 + x_1)$, $|y - x| \leq |y - \tilde{x}|$, we arrive at

$$|(1)| \lesssim \frac{x_1}{|y - \bar{x}||y - \tilde{x}||y - x|^2}.$$

Write $\gamma = \gamma_1 + \gamma_2$ and noting that $|y - \bar{x}| \geq x_2^\gamma |y - \bar{x}|^{1-\gamma}$, $|y - \tilde{x}| \geq x_1^{1-\gamma_2} |y - \tilde{x}|^{\gamma_2}$ and the reflection relations $|y - \tilde{x}|, |y - \bar{x}| \geq |y - x|$ for $y \in [0, 1]^2$, we arrive at

$$|(1)| \lesssim x_2^{-\gamma} x_1^{\gamma_2} |y - x|^{-(3-\gamma+\gamma_2)}.$$

To estimate (2), we use the relation

$$|y + x|^2(y_1 - x_1)^2 - |y - \bar{x}|^2(y_1 + x_1)^2 = -4x_1y_1(y_2 + x_2)^2$$

and similar estimations as above to arrive at

$$\begin{aligned} |(2)| &\lesssim \frac{y_2y_1x_1(y_2 + x_2)^2}{|y - \tilde{x}|^4|y - \bar{x}|^2|y + x|^2} \lesssim \frac{x_1}{|y - \bar{x}||y - \tilde{x}|^3} \\ &\lesssim x_2^{-\gamma} x_1^{\gamma_2} |y - x|^{-3-\gamma_2+\gamma}. \end{aligned}$$

(3)-(6) is mutatis mutandis the same. □

Figure 4 illustrates the domains we need in the proof of the following propositions.

Proposition 9.5. *Let $I = B(x, \frac{1}{2}d_1(x)) \cap \widehat{D}$. Then*

$$\left| P.V. \int_I \frac{\partial(G_i^1 + G_i^2)}{\partial x_j} \omega(y) dy \right| \lesssim Mx_i^{-\alpha} (1 + \delta_{j2} |\log d(x)|)$$




FIGURE 4. Domains of integration in Proposition 9.5.

Proof. Let $0 < \delta < \frac{1}{2}d_1(x)$ so small such that $B(x, \delta) \subset \widehat{D}$. By Proposition 9.2,

$$(70) \quad \left| \int_{I \setminus B(x, \delta)} \sum_{k=1,2} \frac{\partial G_i^k}{\partial x_j} \omega(y) dy \right| \lesssim \left| \int_{I \setminus B(x, \delta)} \sum_{k=1,2} \frac{\partial G_i^k}{\partial y_j} \omega(y) dy \right| + \left| x_i^{-2} \int_{I \setminus B(x, \delta)} \mathcal{O}(|y - x|^{-1}) \right|$$

We distinguish the cases $i = j$ and $i \neq j$. First let $i = j$. Integration by parts gives

$$(70) \lesssim \left| \int_{I \setminus B(x, \delta)} \sum_{k=1,2} G_i^k \frac{\partial \omega}{\partial y_i}(y) dy \right| + \left| x_i^{-2} \int_{I \setminus B(x, \delta)} \mathcal{O}(|y - x|^{-1}) \right| \\ + \sum_{k=1,2} \left(\left| \int_{\partial I} G_i^k \omega(y) \nu_i d\sigma \right| + \left| \int_{\partial B(x, \delta)} G_i^k \omega(y) \nu_i d\sigma \right| \right).$$

where $\nu = (\nu_1, \nu_2)$ is the unit outward pointing normal on ∂I . We first take care of the integral over ∂I . Observe that for $x \in D$, ∂I is either a full circle or the union of a part of a circle and a flat part Σ . Hence

$$\int_{\partial I} |G_i^k \nu_j| d\sigma \leq \int_{\Sigma} |G_i^k| \delta_{2i} d\sigma + \int_{\partial B(0, \frac{1}{2}d_1(x))} |G_i^k| d\sigma$$

For all sufficiently small $\varepsilon > 0$,

$$\begin{aligned}
\int_{\Sigma} |G_i^k| d\sigma &\leq \int_{\Sigma \cap \{|y_1 - x_1| \leq \varepsilon\}} |G_i^k| dy_1 + \int_{\Sigma \cap \{|y_1 - x_1| \geq \varepsilon\}} |G_i^k| dy_1 \\
&\lesssim \int_{\Sigma \cap \{|y_1 - x_1| \leq \varepsilon\}} \frac{x_i^{-1}}{|y - x|} dy_1 + \int_{\Sigma \cap \{|y_1 - x_1| \geq \varepsilon\}} \frac{x_i^{-1}}{|y - x|} dy_1 \\
&\lesssim x_i^{-1} \int_{x_1 - \varepsilon}^{x_1 + \varepsilon} \frac{1}{|x_2 - \delta_2|} dy_1 + x_i^{-1} \int_{1 > |x_1 - y_1| > \varepsilon} \frac{1}{|y_1 - x_1|} dy_1 \\
&\lesssim \frac{x_i^{-1} \varepsilon}{|x_2 - \delta_2|} + x_i^{-1} \int_{\varepsilon}^1 \frac{1}{y_1} dy_1 \\
&\lesssim \frac{x_i^{-1} \varepsilon}{|x_2 - \delta_2|} + x_i^{-1} |\log \varepsilon|.
\end{aligned}$$

Here we used proposition 9.1 again. Choosing $\varepsilon = |x_2 - \delta_2| = d(x)$ we get

$$\int_{\Sigma} |G_i^k| d\sigma \lesssim x_i^{-1} (1 + |\log d(x)|).$$

The other part is estimated by (using proposition 9.1 again)

$$\int_{\partial B(0, \frac{1}{2}d_1(x))} |G_i^k| d\sigma \lesssim x_i^{-1} \int_0^{2\pi} |y - x|^{-1} d_1(x) d\varphi \lesssim x_i^{-1}.$$

Therefore we get for the integral over ∂I , using (69) and (68):

$$\begin{aligned}
\left| \int_{\partial I} G_i^k \omega(y) \nu_i d\sigma \right| &\lesssim \int_{\partial I} |G_i^k \nu_i| |\omega(y)| d\sigma \lesssim M \int_{\partial I} |G_i^k \nu_i| y_i^{1-\alpha} d\sigma \\
&\lesssim M x_i^{1-\alpha} \int_{\partial I} |G_i^k \nu_i| d\sigma \lesssim M x_i^{1-\alpha} x_i^{-1} (1 + \delta_{i2} |\log d(x)|).
\end{aligned}$$

Similar estimates yield that the contribution from the integral over $\partial B(x, \delta)$ is $\lesssim M x_i^{-\alpha}$, with universal constants independent of δ . For

$$x_i^{-2} \int_{I \setminus B(x, \delta)} \mathcal{O}(|y - x|^{-1}) |\omega(y)| dy$$

we obtain the upper bound $\lesssim M x_i^{-\alpha}$ by the same methods.

For the remaining integral we use (68):

$$\begin{aligned}
\left| \int_{I \setminus B(x, \delta)} G_i^k \frac{\partial \omega}{\partial y_i}(y) dy \right| &\lesssim M x_i^{-\alpha} \int_{I \setminus B(x, \delta)} |G_i^k| dy \lesssim M x_i^{-\alpha} \int_{I \setminus B(x, \delta)} |y - x|^{-1} x_i^{-1} dy \\
&\lesssim M x_i^{-\alpha} x_i^{-1} \int_{\delta}^{d_1(x)} \frac{1}{\rho} d\rho \lesssim M x_i^{-\alpha} x_i^{-1} \int_{\delta}^{d_1(x)} \frac{1}{\rho} d\rho \\
&\lesssim M x_i^{-\alpha} x_i^{-1} d_1(x).
\end{aligned}$$

Since $d_1(x) \leq x_i$ we get:

$$\left| \int_{I \setminus B(x, \delta)} G_i^k \frac{\partial \omega}{\partial y_j}(y) dy \right| \lesssim M x_i^{-\alpha}.$$

This concludes the case $i = j$.

For the case $i \neq j$, we have to use the cancellation provided by Proposition 9.4 with $\gamma_1 = 2, \gamma_2 = -1$.

$$\begin{aligned}
\left| \int_{I \setminus B(x, \delta)} \frac{\partial(G_i^1 + G_i^2)}{\partial y_j} \omega(y) \right| dy &\lesssim M \int_{I \setminus B(x, \delta)} \left| \frac{\partial(G_i^1 + G_i^2)}{\partial y_j} \right| y_i^{1-\alpha} dy \\
&\lesssim M x_i^{1-\alpha} \int_{I \setminus B(x, \delta)} y_i^{-1} y_j^{-1} |y - x|^{-1} dy \\
&\lesssim M x_i^{1-\alpha} x_i^{-1} x_j^{-1} \int_{\delta}^{d_1(x)} \frac{1}{\rho} \rho d\rho \\
&\lesssim M x_i^{1-\alpha} x_i^{-1} x_j^{-1} d_1(x) \lesssim M x_i^{-\alpha},
\end{aligned}$$

since $d_1(x) \lesssim x_j$. □

Lemma 9.6. *Let $\gamma \in (0, 1)$ and $x_1 \geq 0$. Then*

$$y_1^\gamma \leq |y_1 - x_1|^\gamma + x_1^\gamma \quad (y_1 \geq 0).$$

Proof. If $y_1 \leq x_1$, the inequality is obvious. For $y_1 > x_1$ we have $y_1 \geq y_1 - x_1 > 0$ and hence $\gamma y_1^{\gamma-1} \leq \gamma(y_1 - x_1)^{\gamma-1}$ so that

$$y_1^\gamma - x_1^\gamma \leq \gamma \int_{x_1}^{y_1} s^{\gamma-1} ds \leq \gamma \int_{x_1}^{y_1} (s - x_1)^{\gamma-1} ds = (y_1 - x_1)^\gamma.$$

□

Proposition 9.7. *Let $II = \widehat{D} \setminus I$ with I as in proposition 9.5. Then*

$$\left| \int_{II} \frac{\partial G_i^k}{\partial x_j} \omega(y) dy \right| \lesssim M x_i^{-\alpha}$$

Proof. Here we have to distinguish two cases. Assume first $d_1(x) = x_i$. Then using proposition 9.1, (69) and Lemma 9.6 we have

$$\begin{aligned}
\left| \int_{II} \frac{\partial G_i^k}{\partial x_j} \omega(y) dy \right| &\lesssim \int_{II} \left| \frac{\partial G_i^k}{\partial x_j} \right| |\omega(y)| dy \\
&\lesssim M \int_{II} |y - x|^{-3} y_i^{1-\alpha} dy \lesssim M \int_{II} |y - x|^{-3} (|y_i - x_i|^{1-\alpha} + x_i^{1-\alpha}) \\
&\lesssim M \int_{II} |y - x|^{-2-\alpha} dy + M x_i^{1-\alpha} \int_{II} |y - x|^{-3} \\
&\lesssim M \int_{\frac{1}{2}d_1(x)}^{\infty} \frac{1}{\rho^{2+\alpha}} \rho d\rho + M x_i^{1-\alpha} \int_{\frac{1}{2}d_1(x)}^{\infty} \frac{1}{\rho^3} \rho d\rho \\
&\lesssim M d_1(x)^{-\alpha} + M x_i^{1-\alpha} d_1(x)^{-1} \lesssim M x_i^{-\alpha} + M x_i^{1-\alpha} x_i^{-1} \lesssim M x_i^{-\alpha}.
\end{aligned}$$

Now let $d_1(x) = x_r$, $r \neq i$. Without loss of generality, we write down only the case $i = 1$ (so $d_1(x) = x_2$). From the explicit relations in Proposition 9.10 and the reflection inequalities we get

$$\left| \frac{\partial G_1^k}{\partial x_i} \right| \leq |y - x|^{-2} |y - \tilde{x}|^{-1}$$

and hence again by (69) and Lemma 9.6,

$$\begin{aligned} \left| \int_{II} \frac{\partial G_1^k}{\partial x_i} \omega(y) dy \right| &\lesssim M \int_{II} \left| \frac{\partial G_1^k}{\partial x_i} \right| y_2^{1-\alpha} dy \\ &\lesssim M \int_{II} |y-x|^{-1-\alpha} |y-\tilde{x}|^{-1} dy + M x_2^{1-\alpha} \int_{II} |y-x|^{-2} |y-\tilde{x}|^{-1} \end{aligned}$$

We continue with the integral without the factor $x_2^{1-\alpha}$ in front; first we enlarge the integration domain by replacing II with $\widehat{D}$, then we split the integration domain into a ball $B(0, 2x_1)$ and the rest.

$$\begin{aligned} \int_{\widehat{D}} |y-x|^{-1-\alpha} |y-\tilde{x}|^{-1} dy &\lesssim \int_{\widehat{D} \cap B(0, 2x_1)} |y-x|^{-1-\alpha} |y-\tilde{x}|^{-1} dy \\ &\quad + \int_{\widehat{D} \setminus B(0, 2x_1)} |y-x|^{-1-\alpha} |y-\tilde{x}|^{-1} dy \\ &\lesssim x_1^{-1} \int_{\widehat{D} \cap B(x, 10x_1)} |y-x|^{-1-\alpha} dy + \int_{B(0, 10) \setminus B(0, 2x_1)} |y|^{-2-\alpha} dy \\ &\lesssim x_1 x_1^{1-\alpha} + x_1^{-\alpha} \lesssim x_1^{-\alpha}. \end{aligned}$$

Here, we used that $x_2 \leq x_1$ (since $d_1(x) = x_2$), so that $|y| \lesssim |y-x|$, $|y| \lesssim |y-\tilde{x}|$ in the second integral. Continuing with $x_2^{1-\alpha} \int_{II} |y-x|^{-2} |y-\tilde{x}|^{-1}$, we get

$$\begin{aligned} x_2^{1-\alpha} \int_{II} |y-x|^{-2} |y-\tilde{x}|^{-1} dy &\lesssim x_2^{1-\alpha} x_1^{-\alpha} \int_{II} |y-x|^{-3+\alpha} dy \\ &\lesssim x_2^{1-\alpha} x_1^{-\alpha} x_2^{-1+\alpha} \alpha x_1^{-\alpha} \end{aligned}$$

using $|y-\tilde{x}| \geq x_1^\alpha |y-\tilde{x}|^{1-\alpha} \geq x_1^\alpha |y-x|^{1-\alpha}$. □

Proposition 9.8. *For $i \neq j$,*

$$\left| \int_{[0,1]^2 \setminus \widehat{D}} \left[\frac{\partial G_i^1}{\partial x_j} + \frac{\partial G_i^2}{\partial x_j} \right] \omega(y) dy \right| \leq C(\gamma_1, \gamma_2) x_i^{-(\gamma_1+\gamma_2)} x_j^{\gamma_2} d(x)^{-1+\gamma_1}$$

where $\gamma_1 \in (0, 1)$, $\gamma_2 \in [0, 1)$, $\gamma_1 + \gamma_2 < 1$. Also,

$$\left| \int_{[0,1]^2 \setminus \widehat{D}} \left[\frac{\partial G_i^1}{\partial x_1} + \frac{\partial G_i^2}{\partial x_1} \right] \omega(y) dy \right| \leq C(\gamma_1) x_i^{-\gamma_1} d(x)^{-1+\gamma_1}$$

Proof. As a preparation, we note that for $x \in D$, $0 < \gamma_1 < 1$,

$$(71) \quad \int_{[0,1]^2 \setminus \widehat{D}} |y-x|^{-3+\gamma_1} \lesssim d(x)^{-1+\gamma_1}.$$

This follows from

$$\begin{aligned} \int_{[0,1]^2 \setminus \widehat{D}} |y-x|^{-3+\gamma_1} &\leq \int_{[0,1]^2 \setminus B(x, d(x))} |y-x|^{-3+\gamma_1} \\ &\leq \int_{B(x, 10) \setminus B(x, d(x))} |y-x|^{-3+\gamma_1}, \end{aligned}$$

since $[0, 1]^2 \setminus \widehat{D}$ is contained in $[0, 1]^2 \setminus B(x, d(x))$ because of $\delta_2 < \delta_3$ and $\delta_1 < \delta_2$.

From Proposition 9.4 we get in case $i \neq j$

$$\begin{aligned} \left| \int_{[0,1]^2 \setminus \widehat{D}} \left[\frac{\partial G_i^1}{\partial x_1} + \frac{\partial G_i^2}{\partial x_j} \right] \omega(y) dy \right| &\leq x_i^{-(\gamma_1+\gamma_2)} x_j^{\gamma_2} \int_{[0,1]^2 \setminus \widehat{D}} |y-x|^{-3+\gamma_1} \\ &\leq x_i^{-(\gamma_1+\gamma_2)} x_j^{\gamma_2} d(x)^{-1+\gamma_1}, \end{aligned}$$

according to (71).

For the second inequality of the Proposition, we note that

$$\left| \frac{\partial G_i^k}{\partial x_i} \right| \lesssim x_i^{-\gamma_1} |y-x|^{-3+\gamma_1},$$

and use (71). □

Proposition 9.9. *For $x \in D$,*

$$\begin{aligned} \left| P.V. \int_{[0,1]^2} \left[\frac{\partial G_i^1}{\partial x_j} + \frac{\partial G_i^2}{\partial x_j} \right] \omega(y) dy \right| &\leq M x_i^{-\alpha} (1 + \delta_{j2} |\log d(x)|) \\ &\quad + C(\gamma_1, \gamma_2) x_i^{-(\gamma_1+\gamma_2)} x_j^{\gamma_2} d(x)^{-1+\gamma_1} \quad (i \neq j) \\ \left| P.V. \int_{[0,1]^2} \left[\frac{\partial G_i^1}{\partial x_i} + \frac{\partial G_i^2}{\partial x_i} \right] \omega(y) dy \right| &\leq M x_i^{-\alpha} (1 + \delta_{i2} |\log d(x)|) + C(\gamma_1) x_i^{-\gamma_1} d(x)^{-1+\gamma_1} \end{aligned}$$

with $\gamma, \gamma_1 \in (0, 1), \gamma_2 \in [0, 1), \gamma_1 + \gamma_2 < 1$.

Proof. We split the integral into a principal value integral over $\widehat{D}$ and a convergent integral over $[0, 1]^2 \setminus \widehat{D}$. The integral over $\widehat{D}$ is further split in to integrals over the domains $I = B(x, \frac{1}{2}d_1(x))$ and $II = \widehat{D} \setminus I$, which are estimated by Propositions 9.5 and 9.7. The part over $[0, 1]^2 \setminus \widehat{D}$ is estimated by Proposition 9.8. □

9.2. Appendix B.

Derivation of Q_i and G_i^j .

In all integrals over infinite domains it is understood that ω is extended periodically and the integrals are understood as limits in the mean. We derive only Q_1 and the formulas for G_1^1, G_1^2, Q_2, G_2^1 and G_2^2 are analogous. The first component of the velocity field is

$$\begin{aligned} u_1(x, t) &= \frac{1}{2\pi} \int_{\mathbb{R}^2} \frac{-(y_2 - x_2)}{|y - x|^2} \omega(y, t) dy \\ &= \frac{1}{2\pi} \left\{ \int_{(0, \infty)^2} + \int_{(-\infty, 0) \times (0, \infty)} + \int_{(0, \infty) \times (-\infty, 0)} + \int_{(-\infty, 0)^2} \right\} \frac{-(y_2 - x_2)}{|y - x|^2} \omega(y, t) dy. \end{aligned}$$

Recall $\tilde{x} = (-x_1, x_2)$, $\bar{x} = (x_1, -x_2)$. Using the double-odd symmetry of ω we can write

$$\begin{aligned} \int_{(-\infty, 0) \times (0, \infty)} \frac{-(y_2 - x_2)}{|y - x|^2} \omega(y, t) \, dy &= \int_{(0, \infty)^2} \frac{y_2 - x_2}{|y - \tilde{x}|^2} \omega(y, t) \, dy \\ \int_{(0, \infty) \times (-\infty, 0)} \frac{-(y_2 - x_2)}{|y - x|^2} \omega(y, t) \, dy &= \int_{(0, \infty)^2} \frac{-(y_2 + x_2)}{|y - \bar{x}|^2} \omega(y, t) \, dy \\ \int_{(-\infty, 0)^2} \frac{-(y_2 - x_2)}{|y - x|^2} \omega(y, t) \, dy &= \int_{(0, \infty)^2} \frac{y_2 + x_2}{|y + x|^2} \omega(y, t) \, dy. \end{aligned}$$

Next we group the integrals in the following way

$$\begin{aligned} &\int_{(0, \infty)^2} \frac{-(y_2 - x_2)}{|y - x|^2} \omega(y, t) \, dy + \int_{(0, \infty)^2} \frac{y_2 - x_2}{|y - \tilde{x}|^2} \omega(y, t) \, dy \\ &= -4x_1 \int_{(0, \infty)^2} \underbrace{\frac{y_1(y_2 - x_2)}{|y - x|^2 |y - \tilde{x}|^2}}_{G_1^1(x, y)} \omega(y, t) \, dy \\ &\quad \int_{(0, \infty)^2} \frac{-(y_2 + x_2)}{|y - \bar{x}|^2} \omega(y, t) \, dy + \int_{(0, \infty)^2} \frac{y_2 + x_2}{|y + x|^2} \omega(y, t) \, dy \\ &= -4x_1 \int_{(0, \infty)^2} \underbrace{\frac{y_1(y_2 + x_2)}{|y + x|^2 |y - \bar{x}|^2}}_{G_1^2(x, y)} \omega(y, t) \, dy. \end{aligned}$$

Together we obtain

$$\begin{aligned} u_1(x, y) &= -x_1 \frac{2}{\pi} \int_{[0, 1]^2} [G_1^1(x, y) + G_1^2(x, y)] \omega(y, t) \, dy \\ &\quad - x_1 \frac{2}{\pi} \int_{\mathbb{R}_+^2 \setminus [0, 1]^2} [G_1^1(x, y) + G_1^2(x, y)] \omega(y, t) \, dy \\ &= -x_1 \left(\frac{2}{\pi} \int_{[0, 1]^2} [G_1^1(x, y) + G_1^2(x, y)] \omega(y, t) \, dy + Q_1^r(x, t) \right) = -x_1 Q_1(x, t). \end{aligned}$$

Proposition 9.10. *The following relations hold:*

$$\begin{aligned} \frac{\partial G_1^1}{\partial x_1} &= -\frac{2y_1(y_1 + x_1)(y_2 - x_2)}{|y - x|^2 |y - \tilde{x}|^4} + \frac{2y_1(y_1 - x_1)(y_2 - x_2)}{|y - x|^4 |y - \tilde{x}|^2} \\ \frac{\partial G_1^1}{\partial x_2} &= \frac{2y_1(y_2 - x_2)^2}{|y - x|^2 |y - \tilde{x}|^4} + \frac{2y_1(y_2 - x_2)^2}{|y - x|^4 |y - \tilde{x}|^2} - \frac{y_1}{|y - x|^2 |y - \tilde{x}|^2} \\ \frac{\partial G_1^1}{\partial y_1} &= -\frac{2y_1(y_1 + x_1)(y_2 - x_2)}{|y - x|^2 |y - \tilde{x}|^4} - \frac{2y_1(y_1 - x_1)(y_2 - x_2)}{|y - x|^4 |y - \tilde{x}|^2} + \frac{y_2 - x_2}{|y - x|^2 |y - \tilde{x}|^2} \\ \frac{\partial G_1^1}{\partial y_2} &= -\frac{2y_1(y_2 - x_2)^2}{|y - x|^2 |y - \tilde{x}|^4} - \frac{2y_1(y_2 - x_2)^2}{|y - x|^4 |y - \tilde{x}|^2} + \frac{y_1}{|y - x|^2 |y - \tilde{x}|^2} \end{aligned}$$

$$\begin{aligned}
\frac{\partial G_1^2}{\partial x_1} &= -\frac{2y_1(y_1+x_1)(y_2+x_2)}{|y+x|^4|y-\bar{x}|^2} + \frac{2y_1(y_1-x_1)(y_2+x_2)}{|y+x|^2|y-\bar{x}|^4} \\
\frac{\partial G_1^2}{\partial x_2} &= -\frac{2y_1(y_2+x_2)^2}{|y+x|^4|y-\bar{x}|^2} - \frac{2y_1(y_2+x_2)^2}{|y+x|^2|y-\bar{x}|^4} + \frac{y_1}{|y+x|^2|y-\bar{x}|^2} \\
\frac{\partial G_1^2}{\partial y_1} &= -\frac{2y_1(y_1+x_1)(y_2+x_2)}{|y+x|^4|y-\bar{x}|^2} - \frac{2y_1(y_1-x_1)(y_2+x_2)}{|y+x|^2|y-\bar{x}|^4} + \frac{y_2+x_2}{|y+x|^2|y-\bar{x}|^2} \\
\frac{\partial G_1^2}{\partial y_2} &= -\frac{2y_1(y_2+x_2)^2}{|y+x|^4|y-\bar{x}|^2} - \frac{2y_1(y_2+x_2)^2}{|y+x|^2|y-\bar{x}|^4} + \frac{y_1}{|y+x|^2|y-\bar{x}|^2} \\
\\
\frac{\partial G_2^1}{\partial x_1} &= -\frac{2y_2(y_1+x_1)^2}{|y+x|^4|y-\tilde{x}|^2} - \frac{2y_2(y_1+x_1)^2}{|y+x|^2|y-\tilde{x}|^4} + \frac{y_2}{|y+x|^2|y-\tilde{x}|^2} \\
\frac{\partial G_2^1}{\partial x_2} &= -\frac{2y_2(y_1+x_1)(y_2+x_2)}{|y+x|^4|y-\tilde{x}|^2} + \frac{2y_2(y_1+x_1)(y_2-x_2)}{|y+x|^2|y-\tilde{x}|^4} \\
\frac{\partial G_2^1}{\partial y_1} &= -\frac{2y_2(y_1+x_1)^2}{|y+x|^4|y-\tilde{x}|^2} - \frac{2y_2(y_1+x_1)^2}{|y+x|^2|y-\tilde{x}|^4} + \frac{y_2}{|y+x|^2|y-\tilde{x}|^2} \\
\frac{\partial G_2^1}{\partial y_2} &= -\frac{2y_2(y_1+x_1)(y_2+x_2)}{|y+x|^4|y-\tilde{x}|^2} - \frac{2y_2(y_1+x_1)(y_2-x_2)}{|y+x|^2|y-\tilde{x}|^4} + \frac{y_1+x_1}{|y+x|^2|y-\tilde{x}|^2} \\
\\
\frac{\partial G_2^2}{\partial x_1} &= \frac{2y_2(y_1-x_1)^2}{|y-x|^2|y-\bar{x}|^4} + \frac{2y_2(y_1-x_1)^2}{|y-x|^4|y-\bar{x}|^2} - \frac{y_2}{|y-x|^2|y-\bar{x}|^2}, \\
\frac{\partial G_2^2}{\partial x_2} &= -\frac{2y_2(y_1-x_1)(y_2+x_2)}{|y-x|^2|y-\bar{x}|^4} + \frac{2y_2(y_1-x_1)(y_2-x_2)}{|y-x|^4|y-\bar{x}|^2}, \\
\frac{\partial G_2^2}{\partial y_1} &= -\frac{2y_2(y_1-x_1)^2}{|y-x|^2|y-\bar{x}|^4} - \frac{2y_2(y_1-x_1)^2}{|y-x|^4|y-\bar{x}|^2} + \frac{y_2}{|y-x|^2|y-\bar{x}|^2}, \\
\frac{\partial G_2^2}{\partial y_2} &= -\frac{2y_2(y_1-x_1)(y_2+x_2)}{|y-x|^2|y-\bar{x}|^4} - \frac{2y_2(y_1-x_1)(y_2-x_2)}{|y-x|^4|y-\bar{x}|^2} + \frac{y_1-x_1}{|y-x|^2|y-\bar{x}|^2}.
\end{aligned}$$

REFERENCES

1. H. Bahouri, J.-Y. Chemin, *Equations de transport relatives a des champs de vecteurs non-Lipschitziens et mecanique de fluides*, (French) Arch. Rational Mech. Anal. 127, **2**, 159-181 (1994).
2. D. Córdoba: *Nonexistence of Simple Hyperbolic Blow-Up for the Quasi-Geostrophic Equation*, Ann. Math., Second Series, Vol. 148, No. 3 (1998), 1135-1152
3. D. Córdoba, C. Fefferman: *Growth of Solutions for QG and 2D Euler Equations*, Journal of the American Mathematical Society, **15**, 3, p. 665-670 (2002).
4. S. Denisov: *Infinite superlinear growth of the gradient for the two-dimensional Euler equation*, Discrete Contin. Dyn. Syst. A 23 (2009), 755-764.
5. S. Denisov: *Double-exponential growth of the vorticity gradient for the two-dimensional Euler equation*, Proceedings of the AMS, Vol. 143, N3, 2015, 1199-1210.
6. S. Denisov: *The sharp corner formation in 2D Euler dynamics of patches: infinite double-exponential rate of merging*, Arch. Rational Mech. Anal., Vol. 215, N2, 2015, 675-705.
7. N. H. Katz and A. Tapay, *A model for studying double exponential growth in the two-dimensional Euler equations*, preprint [arXiv:1403.6867](https://arxiv.org/abs/1403.6867).
8. A. Kiselev and V. Šverák, *Small scale creation for solutions of the incompressible two dimensional Euler equation*, Annals of Math. 180 (2014), 12051220.
9. W. Walter, *Differential and Integral Inequalities*, Springer 1970.

10. D. Wilett, *A Linear Generalization of Gronwall's inequality*, Trans. Am. Math. Soc, **16**, 774-778 (1965).
11. A. Zlatoš, *Exponential growth of the vorticity gradient for the Euler equation on the torus*, Adv. Math. 268 (2015), 396-403.

RICE UNIVERSITY, DEPARTMENT OF MATHEMATICS-MS 136, Box 1892, HOUSTON, TX 77251-1892

E-mail address: `hoang@math.wisc.edu`

RICE UNIVERSITY, DEPARTMENT OF MATHEMATICS-MS 136, Box 1892, HOUSTON, TX 77251-1892/INSTITUTE FOR ANALYSIS, KARLSRUHE INSTITUTE FOR TECHNOLOGY (KIT), KAISERSTRASSE 89, 76133 KARLSRUHE (GERMANY)

E-mail address: `radosz@math.uni-karlsruhe.de`